



Understanding Strategic Coastal Blue Carbon Opportunities in the Seas of East Asia

Report by
Partnerships in Environmental Management for the Seas of
East Asia (PEMSEA), Conservation International and
The Nature Conservancy

Report Authors:
Silvestrum Climate Associates, LLC.



Understanding Strategic Coastal Blue Carbon Opportunities in the Seas of East Asia

Report by
Partnerships in Environmental Management for the Seas of
East Asia (PEMSEA), Conservation International and
The Nature Conservancy

Report Authors:
Silvestrum Climate Associates, LLC.



Understanding Strategic Coastal Blue Carbon Opportunities in the Seas of East Asia

May 2017

This publication may be reproduced in whole or in part and in any form for educational or non-profit purposes or to provide wider dissemination for public response, provided prior written permission is obtained from the PEMSEA Resource Facility Executive Director, acknowledgment of the source is made and no commercial usage or sale of the material occurs. PEMSEA would appreciate receiving a copy of any publication that uses this publication as a source.

No use of this publication may be made for resale, any commercial purpose or any purpose other than those given above without a written agreement between PEMSEA and the requesting party.

Crooks, S., von Unger, M., Schile, L., Allen, C. & Whisnant, R. (2017) Understanding Strategic Blue Carbon Opportunities in the Seas of East Asia. Report by Silvestrum Climate Associates for Partnerships in Environmental Management for the Seas of East Asia (PEMSEA), Conservation International and The Nature Conservancy, with support from the Global Environment Facility and United Nations Development Program.

ISBN 978-971-812-040-8

The contents of this publication do not necessarily reflect the views or policies of PEMSEA Country Partners and its other participating organizations, Conservation International or The Nature Conservancy. The designation employed and the presentation do not imply expression of opinion, whatsoever on the part of PEMSEA concerning the legal status of any country or territory, or its authority or concerning the delimitation of its boundaries.

Published by

Partnerships in Environmental Management for the Seas of East Asia (PEMSEA)

P.O. Box 2502, Quezon City 1165, Philippines

Tel: (+632) 929-2992 Fax: (+632) 926-9712

Email: info@pemsea.org

www.pemsea.org

Partnerships in Environmental Management for the Seas of East Asia (PEMSEA) is an intergovernmental organization operating in East Asia to foster and sustain healthy and resilient oceans, coasts, communities and economies through integrated management solutions and partnerships. For over two decades, PEMSEA has provided solutions for effective management of coasts and oceans across the shared seas of East Asia. As the regional coordinating mechanism for the Sustainable Development Strategy for the Seas of East Asia (SDS-SEA), a shared marine strategy among 14 countries in the region, PEMSEA works with national and local governments, companies, research and science institutions, communities, international agencies, regional programs, investors and donors towards implementation of the SDS-SEA.

Conservation International (CI) is a nonprofit environmental organization with the goal to protect nature as a source of food, fresh water, livelihoods and a stable climate. For nearly 30 years, CI has been protecting nature for the benefit of all through science, policy and partnerships with countries, communities and companies. CI works with more than 2,000 partners in 30 countries and has helped support 1,200 protected areas and interventions across 77 countries, safeguarding more than 601 million hectares of land, marine and coastal areas.

The Nature Conservancy is a conservation organization working around the world to protect the lands and waters on which all life depends. With more than 1 million members and 600 scientists, the Conservancy has protected 120 million acres of land and 5,000 miles of rivers worldwide and operates more than 100 marine conservation projects globally.

Silvestrum Climate Associates is a consultancy providing technical analysts and practitioners in the fields of environmental science and policy, coastal engineering and carbon project development, with a mission to assist public and private sector clients tackle challenges of climate change and sustainable development.

The International Blue Carbon Initiative is a coordinated, global program focused on mitigating climate change through the conservation and restoration of coastal and marine ecosystems. The Initiative comprises a global network of scientists, practitioners and policy advisors collaborating to advance improved management of blue carbon ecosystems.

Acknowledgements

The authors of this report thank the World Conservation Monitoring Center, Dr. Daniel Friess at the National University of Singapore and The Nature Conservancy for sharing data files that supported this analysis. Dr. Miguel Fortes of the University of the Philippines provided helpful comments on regional seagrass status and trends. We thank also Dr. Emily Pidgeon of Conservation International and Emily Landis of The Nature Conservancy for review of document drafts.

Table of Contents

Acronyms and Abbreviations	v
Executive Summary	1
I. Introduction	5
II. Status of Coastal Blue Carbon Ecosystems in East Asia	7
Mangrove Ecosystems	7
Tidal Marsh Ecosystems	11
Seagrass Ecosystems	13
Historic Loss of Blue Carbon Ecosystems	15
III. Blue Carbon as a Climate Intervention Concept: UNFCCC, Kyoto Protocol and the Paris Agreement	17
History I: Between Mitigation and Adaptation	18
History II: Blue Carbon Accounting Rules and Practice	18
CDM: Mangrove Afforestation and Reforestation Approaches	19
Voluntary Carbon Markets	19
REDD: Overlapping Concepts	20
Blue Carbon and NAMAs	21
IV. Implementing Blue Carbon Under Nationally Determined Contributions (NDCs)	23
NDC Commitments by Countries in East Asia	23
Moving Forward Under NDCs in the Region	28
Policy Opportunities for PEMSEA Partner Countries	30
Linking Climate Finance and Blue Carbon Investment	33
V. Practical Steps to Advance Blue Carbon Interventions	35
VI. Addendum: Blue Carbon Ecosystems as Part of an East Asia Blue Economy	39
Private Investment for Conserving and Restoring Blue Carbon Ecosystems	39
Blue Carbon and the Blue Economy	41
VII. Appendices	43
Appendix A: Definitions	43
Appendix B: Maps of Mangrove Extent and Tidal Wetland Losses	45
Appendix C: Bibliography	53

Acronyms and Abbreviations

CDM	Clean Development Mechanism
CER	Certified Emission Reduction
CI	Conservation International
FLEGT+	Forest Law Enforcement Governance and Trade
GHG	Greenhouse Gas
ha	Hectare
ICM	Integrated Coastal Management
IPCC	Intergovernmental Panel on Climate Change
JCM	Joint Crediting Mechanism
LULUCF	Land Use, Land-Use Change and Forestry
MoU	Memorandum of Understanding
Mha	Million Hectares
MMt C	Million Metric Tons of Carbon
MMt CO₂	Million Metric Tons of Carbon Dioxide
MPA	Marine Protected Area
NAMA	National Appropriate Mitigation Action
NC	National Communication
NDC	Nationally Determined Contribution
OECD	Organisation for Economic Co-operation and Development
PEMSEA	Partnerships in Environmental Management for the Seas of East Asia
REDD+	Reducing Emissions from Deforestation and forest Degradation, the Conservation of Forest Carbon Stocks, Sustainable Management of Forests, and Enhancement of Forest Carbon Stocks
SBSTA	“Subsidiary Body for Scientific and Technological Advice” to the UNFCCC
SCA	Silvestrum Climate Associates LLC
SDGs	Sustainable Development Goals
SDS-SEA	Sustainable Development Strategy for the Seas of East Asia
TNC	The Nature Conservancy
UNEP	United Nations Environment Programme
UNFCCC	United Nations Framework Convention on Climate Change
VCS	Verified Carbon Standard





Photo by SCA/Crooks

Executive Summary

Coastal blue carbon is a term that recognizes the role of coastal wetlands in the global carbon cycle. Mangroves, tidal marshes and seagrass meadows (collectively called coastal blue carbon ecosystems) sequester carbon dioxide from the atmosphere continuously over thousands of years, building stocks of carbon in biomass and organic rich soils. This ecosystem service is in addition to other ecosystem services provided by coastal wetlands that underpin fish stocks, maintain food security and contribute to filtration of sediment, protecting coral reefs and populated coastal lowlands from erosion and flooding.¹

Despite their importance, coastal wetlands are some of the most threatened ecosystems on Earth, with up to 800,000 hectares destroyed each year, approximately 1.5% of global coverage. When degraded or destroyed, the services provided by these coastal blue carbon ecosystems are not only lost, but notably, they can become a significant source of greenhouse gas emissions, with thousands of years of sequestered carbon released over a period of years to decades.² Although the combined global area of mangroves, tidal marshes and seagrass meadows equates to only 2-6% of the total area of terrestrial tropical forest, its ongoing losses accounts for up to 10-20% of the emissions from global deforestation—a total of 0.5 billion tons of carbon dioxide emissions annually.³

Over the past 10 years, researchers, policymakers and practitioners have built a strong foundation of science, policy, finance and coastal management approaches for integrating the conservation and restoration of blue carbon ecosystems into the global effort to address climate change. Such efforts reflect a growing awareness of the importance of coastal blue carbon ecosystems in terms of global climate regulation and adaptation for local communities. Protecting and restoring blue carbon ecosystems are a key link to achieving sustainable development goals, growing blue economy and meeting national commitments to the Paris Climate Agreement.

East Asia is a global hotspot for remaining coastal blue carbon ecosystems, but it is experiencing high rates of loss. Many East Asian countries are also home to some of the most vulnerable coastal communities to the impacts of climate change and declining coastal ecosystem services. There is an opportunity to improve management of coastal blue carbon ecosystems towards achieving climate change commitments, sustainable development goals and the well-being of coastal communities.

This report seeks to better understand the status of coastal blue carbon ecosystems in East Asia and raise awareness of the opportunities to include improved

¹ Costanza, R. et al. 2014. "Changes in the global value of ecosystem services." *Global Environmental Change*, 26:152-158.

² Crooks, S. et al. 2011. "Mitigating climate change through restoration and management of coastal wetlands and near-shore marine ecosystems: challenges and opportunities." *Environment Department Paper 121*. World Bank, Washington, DC.

³ Pendleton, L. et al. 2012. "Estimating global "blue carbon" emissions from conversion and degradation of vegetated coastal ecosystems." *PloS One*, 7.9:e43542.

management of these ecosystems within climate mitigation and adaptation actions and commitments, including opportunities to access new forms of financing. Its geographic focus is on countries with coastal ecosystems that have signed the Sustainable Development Strategy for the Seas of East Asia (SDS-SEA), specifically: Brunei Darussalam, Cambodia, China, DPR Korea, Indonesia, Japan, Malaysia, Philippines, RO Korea, Singapore, Thailand, Timor-Leste and Viet Nam.

All three blue carbon ecosystems—mangroves, tidal marshes and seagrass meadows—occur in East Asia. Mangroves are the dominant tidal wetland ecosystem type at latitudes between approximately 30°N and 30°S, covering most of East Asia. RO Korea or DPR Korea are the only two countries in this study with no mangrove cover due to their northern extent. Tidal marshes are found at latitudes that are too cold for mangroves to tolerate, typically latitudes greater than 30°N and 30°S. Therefore, in East Asia, tidal marshes occur primarily in China, Japan, DPR Korea and RO Korea. Seagrasses, though extensive, remain very poorly mapped with estimates of extent and rates of loss being derived from experts with knowledge of field surveys and reports.

In all, there are 4 million ha of mangroves in the region, representing around 30% of the global total. Five of the top 11 mangrove-holding countries globally are in East Asia: Indonesia (2.71 Mha), Malaysia (0.56 Mha), the Philippines (0.26 Mha), Thailand (0.25 Mha) and Viet Nam (0.22 Mha). Applying regional forestry models and IPCC default values for soil carbon stocks, it is estimated that the region's mangroves hold 8.8 billion tons of CO₂ vulnerable to release by human activities. Each year, these mangroves remove approximately, 22.4 MMt CO₂ from the atmosphere.

The full extent of tidal marshes is not known. China held 57,344 ha in 1995,⁴ perhaps representing less than 10% of what once existed across temperate areas of East Asia. Remaining vulnerable tidal wetland carbon stocks are estimated to be between 180-660 MMt CO₂. Each year, these wetlands sequester 0.19 MMt CO₂.

It is estimated that there may be 3 million ha of seagrasses across the region, holding 1.3 billion tons CO₂ within soils and sequestering 4.7 MMt CO₂ each year. The lack of data on distribution and carbon stocks in tidal wetland and seagrass ecosystems are critical data gaps that have the potential to be filled. Efforts are underway in the Philippines and Indonesia to quantify and model seagrass carbon stocks and fluxes through the BlueCARES project.

The historic extent of all blue carbon ecosystem area now converted to diked coastal floodplain is calculated to be 3.36 million ha across the region. The five countries with the greatest extent of these subsided coastal lands are China (1.81 Mha), Viet Nam (0.85 Mha), the Philippines (0.68 Mha), Indonesia (0.30 Mha) and Japan (0.23 Mha). As a first approximation, using IPCC default values for soil carbon loss, it is estimated that some 3.7 billion tons CO₂ have emitted from converted mangroves and tidal marshes across the region, and annual carbon sequestration of around 6 MMt CO₂ has been lost. These emissions are likely to be underestimates, since areas where drained organic soil remain, emissions will be continuing to this day.⁵

While losses of tidal marshes and seagrasses have not been assessed, over the period 2000 to 2012 the greatest rates of mangrove loss are found in Malaysia, Cambodia, Indonesia, Thailand and China. The primary causes for mangrove loss are conversion to aquaculture (33,721 ha), oil palm plantations (18,456 ha),

⁴ Shi-Jun, Y. and C. Ji-yu. 1995. "Coastal salt marshes and mangrove swamps in China." *Chinese Journal of Oceanology and Limnology*, 13: 318-324.

⁵ Crooks, S. et al. 2011. "Mitigating climate change through restoration and management of coastal wetlands and near-shore marine ecosystems: challenges and opportunities."

deforestation (5,483 ha) and urbanization (4,476 ha). Across the region, some 17,496 ha of new mangrove growth was identified, representing potential building of new mangrove in delta regions and reforestation on abandoned lands.

The potential to restore blue carbon ecosystems depends upon site specific consideration. Some sites are readily restorable; for others, recovering from former conditions is not possible due to the degree of environmental change.⁶ While the needed experience and expertise to assess restoration potential and enact recovery exists, in most cases, failure is typically due to poor site selection and planning. Restoration of seagrass meadows can often be more challenging than tidal marshes or mangroves though, again, there is an established scientific and restoration practice.

Several opportunities exist for countries to improve management of coastal blue carbon ecosystems. Many countries, including PEMSEA partner countries, recognize the importance of mangroves, if not all blue carbon ecosystems. However, conservation and restoration policies often lack effectiveness. Enforcement capacities are minimal, activities within various government departments are heterogeneous (fisheries as opposed to nature conservation, for instance) and uncertain land tenure hampers the capacity to identify relevant stakeholders, plan with them and secure sustainable investment.

International climate policy offers an opportunity to restate the importance of coastal ecosystems and establish effective mitigation and adaptation tools. The emerging framework of nationally determined contributions (NDC)–country commitments on climate change–creates powerful incentives for action. Almost all PEMSEA countries have started addressing blue carbon ecosystems in their NDCs. While to date, the

mitigation opportunities have often been overlooked, future rounds of NDC restatements will permit greater and more comprehensive focus. This includes an openness for greater bilateral and regional cooperation between economically developed and developing countries; between countries that operate or plan to operate domestic emissions trading systems and those that are willing to generate international offset credits; and between countries that already employ more advanced and comprehensive inventories and others that require capacity building and technological assistance.

Recommendations

There are a number of steps that PEMSEA partner countries can take to build awareness, facilitate knowledge exchange and accelerate practical action on blue carbon. These include:

- 1) Improved tracking of blue carbon ecosystem gains and losses (noting particularly a lack of data on tidal marshes and seagrasses), quantification and reporting of greenhouse emissions and removals following guidance provided in the IPCC Wetland Supplement;⁷
- 2) Including coastal blue carbon ecosystems within NDCs and related policies under national mitigation and adaptation commitments to the Paris Agreement;
- 3) Measuring and weighing the significance of coastal blue carbon ecosystems across policy areas and planning documents, including on trade, aid and integrated coastal management;
- 4) Developing climate change vulnerability assessments, adaptation and resilience plans, and

⁶ *Ibid.*

⁷ Hiraishi, Takahiko, et al. 2014. *2013 supplement to the 2006 IPCC guidelines for national greenhouse gas inventories: Wetlands*. IPCC, Switzerland.

promoting the role of coastal blue carbon ecosystems as a vehicle for sustainable environmental infrastructure;

- 5) Building on bilateral and regional cooperation of PEMSEA countries to work towards joint planning and implementation of the NDC framework;
- 6) Engaging in programs and demonstration activities that build public–private initiatives or support international financing for coastal blue carbon ecosystem conservation and restoration.

While there are actions that each country can take within national borders to support coastal blue carbon ecosystems and their climate benefits, there are additional benefits to be gained through transboundary

collaboration. Sharing of knowledge and experience builds regional capacity (e.g., development of GHG inventories, NDCs, NAMAs, science and restoration experience). Countries advancing carbon finance trade infrastructure may serve as hubs for regional market interventions. Those seeking partners to reduce in-country emissions may engage with other countries requiring support. Donor countries may provide scientific and technical development assistance to regional neighbors. Many of the threats facing coastal blue carbon ecosystems are transboundary in nature (e.g., water quality, sediment delivery, trade impacts) and would benefit from international cooperation to address as part of regional sustainability actions. Regional coordinating mechanisms, such as PEMSEA, can offer institutional framework for enhanced regional planning and implementation.



Photo by SCA/Crooks

1

Introduction

Coasts and oceans are some of the most productive ecosystems on the planet, providing a rich array of ecosystem services that maintain human survival and quality of life, supporting local communities and national economies. More than half of the world's population lives within 200 km of the coast, drawn to environmental and economic resources that support large populations.⁸

Consequently, coastal ecosystems are among the most threatened, under pressure from destruction and degradation. Since the turn of the 19th century, it is estimated that 50% of all mangroves and tidal marshes globally have been converted to other land uses, and area of seagrass meadows have been reduced by 29%.⁹ The loss of these wetlands has resulted in a forfeiture of the beneficial ecosystems services they provided,^{10,11} including food provision, storm protection, climate regulation and local ocean acidification buffering, as well as aesthetic and spiritual benefits.

One key service provided by coastal ecosystems, but historically overlooked, is the regulation of the global climate by coastal ecosystems. The concept of “coastal blue carbon” has emerged to recognize the need for improved management of coastal ecosystems to support this important climate regulating service. Within just a few years, coastal blue carbon has evolved from a mostly scientific interest into a cross-cutting policy tool linking greenhouse gas (GHG) accounting for coastal environments—a major global emissions source—with short- and long-term commitments of a wide range of countries, climate finance (for both mitigation and adaptation) and funding opportunities for developed countries, private investors and coastal communities.

The concept of coastal blue carbon recognizes the role of the ocean's biological systems in buffering the world's atmospheric CO₂ levels. This service is provided through the extraction of CO₂ by plants directly from the atmosphere and surface waters.¹²

⁸ Creel, Liz. 2003. “Ripple Effects: Population and Coastal Regions.” <http://www.prb.org/Publications/Reports/2003/RippleEffectsPopulationandCoastalRegions.aspx>. Accessed 2016 November 3.

⁹ Pendleton, L. et al. 2012. “Estimating global “blue carbon” emissions from conversion and degradation of vegetated coastal ecosystems.”

¹⁰ Costanza, R. et al. 1997. “The value of the world's ecosystem services and natural capital.” *Nature*, 387:253–260.

¹¹ Costanza, R. et al. 2014. “Changes in the global value of ecosystem services.”

¹² Pidgeon, E. 2009. “Carbon sequestration by coastal marine habitats: important missing sinks.” In *The management of natural coastal carbon sinks*. Edited by Laffoley, D. and Grimsditch, G. IUCN, Gland.

Coastal blue carbon is the contribution to climate mitigation made by coastal ecosystems, particularly mangroves, tidal marshes (salt, brackish, and freshwater marshes) and seagrasses.¹³ These ecosystems directly extract CO₂ from the atmosphere and coastal waters and store it as solid form carbon in biomass and soil material. The soil carbon pool is particularly important because low-oxygen conditions created by wet soils significantly decrease decomposition rates, allowing build-up of carbon stocks that exceed those of upland soils.

Approximately 48% of carbon sequestration of the entire ocean occurs within just 2% of its area, through transfer of atmospheric CO₂ to the soil carbon pool.¹⁴ Not recognized in this calculation, but also valuable, is the flux of dissolved and particulate carbon from coastal ecosystems to support food webs in coastal and ocean areas.¹⁵ Some of this “lateral carbon flux” eventually becomes buried in offshore marine sediments, but is difficult to track and not currently included directly in climate mitigation calculations. Thus, while conserving and improving all coastal ecosystems and marine ecological conditions is beneficial to carbon sequestration, blue carbon interventions for meeting quantified GHG mitigation goals are focused on management of mangroves, tidal marshes and seagrasses.

When coastal blue carbon ecosystems are drained or excavated and converted to other land uses, carbon sequestration is halted. More significantly, stored soil carbon stocks are released and, together with the biomass carbon stock (e.g., removed mangrove trees), are returned to the atmosphere. Soil carbon stocks that had accumulated over thousands of years may

be returned within just a few decades.^{16,17,18} It is estimated that some 450 MMt CO₂ of soil carbon from blue carbon ecosystems are released from land use change each year with an annual economic impact of \$18 billion USD.¹⁹ Such levels equate to 10-20% of total CO₂ emissions from global deforestation and rise to a level at which economic instruments may be brought to bear to reduce or reverse environmental losses.²⁰

Blue carbon ecosystems are heavily concentrated in a few countries and regions, including the seas of East Asia, with Southeast Asia as the unambiguous geographic global center. Covering over 7 million km² of sea area and 235,000 km of coastline, East Asia is home to some of the most economically and ecologically important sea areas of the world. With a growing population and ongoing coastal migration, pressure continues to mount on coastal and ocean resources in the region. The environmental and social impact of these activities threatens a decline in coastal ecosystems that form the basis of the economic growth and well-being of the region, and the planet. Many East Asian countries—home to the world’s largest coastal carbon stocks and some of the most vulnerable coastal communities to the impacts of climate change—have embraced the blue carbon concept and are seeking to implement it in the years to come.

This report outlines the status and trends of coastal blue carbon ecosystems in the seas of East Asia and summarizes their relevance to developing climate mitigation commitments, blue carbon interventions and financing options. Finally, recommendations for interventions on policy advancement and direct actions to support management, climate adaptation and development of a regional blue economies are provided.

¹³ Howard, J. et al. 2017. “Clarifying the role of coastal and marine systems in climate mitigation.” *Frontiers in Ecology and the Environment*, 15:42-50.

¹⁴ Duarte, C.M. et al. 2005. “Major role of marine vegetation on the oceanic carbon cycle.” *Biogeosciences*, 2:1–8.

¹⁵ Twilley, R. et al. 1992. “Carbon sinks in mangroves and their implications to carbon budget of tropical coastal ecosystems.” In *Natural Sinks of CO₂*, Springer, Netherlands, pp. 265-288.

¹⁶ Crooks, S. et al. 2011. “Mitigating climate change through restoration and management of coastal wetlands and near-shore marine ecosystems: challenges and opportunities.”

¹⁷ Kauffman, J. B. and D. Donato. 2012. *Protocols for the measurement, monitoring and reporting of structure, biomass and carbon stocks in mangrove forests*. (CIFOR Working Paper no. 86, p. 40p). Center for International Forestry Research (CIFOR), Bogor, Indonesia.

¹⁸ Fourqurean, J.W. et al. 2012. “Seagrass ecosystems as a globally significant carbon stock.” *Nature Geoscience*, 5:505-509.

¹⁹ Pendleton, L. et al. 2012. “Estimating global “blue carbon” emissions from conversion and degradation of vegetated coastal ecosystems.”

²⁰ Murray, B. et al. 2011. *Green payments for blue carbon: Economic incentives for protecting threatened coastal habitats*. Nicholas Institute for Environmental Policy Solutions, Report NI 11.04.



Photo by SCA/Crooks

2

Status of Coastal Blue Carbon Ecosystems in East Asia

Coastal wetlands, which include mangroves, tidal marshes and seagrasses, are found in a narrow elevation range along the coast. Mangroves and tidal marshes occupy the top half of the tidal range, not occurring below mean tide elevations or above the highest tides. Seagrasses are found predominantly below the tides with their lowest depth range determined by water clarity. The following section summarizes current knowledge of each ecosystem's distribution in East Asia, estimated carbon stocks and estimated emissions due to ecosystem loss.

Mangrove Ecosystems

Mangroves are the dominant tidal wetland ecosystem type at latitudes between approximately 30°N and 30°S. This latitudinal range covers most of East Asia. RO Korea or DPR Korea are the only two countries in this study with no mangrove cover due to their northern extent. There are at least 41

mangrove species found within East Asia, making it a significant hotspot for mangrove diversity.

Over the past several years, several efforts have been undertaken to map mangroves globally. In 2000, for instance, four separate approaches were used to assess baseline mangrove cover.²¹ Two estimates were based on mangrove presence/absence in a given area from classified Landsat satellite imagery: Giri et al. (2011)²² and the World Atlas of Mangroves²³ (**Figure 1; Appendix Figures 1-11**). The other estimates were based on mangrove cover within a given area (instead of presence/absence) and were derived from a combination of previous mangrove and forest mapping efforts (Giri et al. 2011 – 'Mangrove Forests of the World' [MFW] and 'Terrestrial Forest of the World' [BIOME]). Mangrove cover was assessed between 2000 and 2012 using imagery and estimated for 2014 based on rates of change.²⁴ None of these methods account for mangrove loss prior to 2000 and thus are only an estimate of recent change.


²¹ In 1999, the Landsat 7 satellite was launched, which captured moderate resolution (30m) imagery across the world at a 16-day interval. In many cases, this was the first imagery of its kind to be collected in many parts of the world and enabled systematic image analysis, such as mapping mangroves, at a global scale.

²² Giri, C. et al. 2011. "Status and distribution of mangrove forests of the world using earth observation satellite data." *Global Ecology and Biogeochemistry*, 20: 154-159.

²³ Spalding, M. 2010. *World atlas of mangroves*. Routledge.

²⁴ Hamilton, S. and D. Casey. 2016. "Creation of a high spatio-temporal resolution global database of continuous mangrove forest cover for the 21st century (CGMFC-21)." *Global Ecology and Biogeochemistry*, 25: 729-738.

Figure 1. Distribution of mangroves in East Asia based on data from the 'World Atlas of Mangroves'. PEMSEA countries with mangroves in medium grey.



Considering that five of the top eleven mangrove-holding countries globally are in East Asia, mangroves are indeed a critical blue carbon ecosystem for the region (**Table 1**). Indonesia has the greatest mangrove coverage of any country, both within East Asia and globally. There are some differences in estimated mangrove coverage depending on the methodology used. Notably, the BIOME method estimates are nearly double all other estimates and are likely an overestimation. This would seem to be the case, based on visual comparisons made by the authors of this report.

Based on the estimated mangrove area, mangrove carbon stocks across the region were calculated using several different approaches. Field collected data provide the most reliable method for quantifying carbon stocks at a given location, and there are challenges to extrapolating these

values across a region as large as East Asia. This can be accomplished by combining multiple data sources to derive a statistical average of soil conditions (for example, the global default values provided by the IPCC 2014) or modelling stocks based on environmental conditions and field data to derive a potential carbon stock.^{25, 26}

For this report, we calculate mangrove biomass and soil carbon stocks from default values and global mangrove modelling to estimate national carbon stocks. Using a field-based carbon stock assessment for Indonesia (where researchers have conducted the most extensive field work in the region), we highlight that national values can vary considerably from the modelled estimates. Hamilton and Friess (in review)⁸ created five global models for estimating biomass (above- and below-ground) and

²⁵ Hamilton, S. E. and D. Friess. 2016. "High-resolution annual maps of whole system mangrove carbon stocks from 2000 to 2012." *arXiv preprint arXiv:1611.00307*.

²⁶ Hutchinson, J. et al. 2013. "Predicting global patterns in mangrove forest biomass." *Conservation Letters*, 7:233-240.

Table 1. Estimated mangrove area in 2000 based on four different techniques, and coverage in 2012 and 2014 based on Hamilton & Casey (2016).

Country	2000				2012		2014 ⁵	
	Giri et al. ^a	MFW ^b	BIOME ^c	WAM ^d	MFW ^b	BIOME ^c	MFW ^b	BIOME ^c
	ha	ha	ha	ha	ha	ha	ha	ha
Indonesia	2,707,572	2,407,313	4,664,152	2,986,392	2,332,429	4,305,957	2,314,277	4,227,806
Malaysia	558,581	496,868	873,795	709,727	472,584	770,043	469,150	761,623
Philippines	259,037	209,105	211,515	257,780	206,424	208,761	205,975	208,450
Thailand	245,121	193,345	436,165	250,057	188,633	399,979	187,562	393,611
Viet Nam	215,529	71,640	128,791	101,497	70,817	126,293	70,641	125,930
Cambodia	47,572	33,839	75,339	60,189	32,322	65,375	32,004	64,078
China	17,910	3,223	3,580	20,118	3,155	3,491	3,149	3,487
Brunei Darussalam	11,089	10,423	14,652	17,134	10,341	14,345	10,327	14,341
Timor-Leste	1,067	857	853	n.d.	846	843	844	842
Japan	1,000	792	803	652	786	797	784	795
Singapore	583	167	167	464	167	165	167	164
Total	4,065,061	3,427,572	6,409,812	4,404,010	3,318,504	5,896,049	3,294,880	5,801,127

^a Giri et al. 2011 dataset; ^b Mangrove Forests of the World dataset (Hamilton and Casey 2016); ^c Terrestrial Ecosystems of the World dataset (Hamilton and Casey 2016); ^d World Atlas of Mangrove dataset (imagery between 1999 & 2003); n.d. = no data

converted these values to carbon using a conversion factor of 47.5%, combining tree carbon stocks with modelled soil carbon stocks to 1 m depth.²⁷ This study averages the results from these models in **Table 2**. In another example, Hutchinson et al. (2013) created a bioclimatic model to estimate above- and below-ground biomass. These values are converted to carbon using the same 47.5% conversion factor. Hutchinson et al. (2013)⁹ restricts the analysis to countries with mangrove coverage >5,000 ha. Therefore, there are no data for Singapore, Japan and Timor-Leste from the Hutchinson study.

Since soil carbon stocks were not determined in this initial analysis, two values from the IPCC soil carbon stock for mangroves down to 1 m depth are applied in this study, one specifically for organic soil and one that is an aggregate of organic and mineral soils.²⁸ Total carbon stocks and CO₂ equivalents (CO₂ eq.) are summarized by country in **Table 2**. In line with mangrove coverage, Indonesia is estimated to

have the highest mangrove carbon stocks, followed by Malaysia and the Philippines. Mangrove carbon stocks in Japan are the lowest due to their limited extent. Since all mangrove carbon stock estimates used in this analysis were based on national estimates derived from global models, these modelled values are compared with in-situ measurements determined by country-specific data. In Indonesia, Alongi et al.²⁹ calculated total mangrove carbon stocks, based on multiple mangrove field surveys and soil cores (standardized to 1 m depth) conducted across the country, to be 3,031 MMt C (or 11,114 MMt CO₂). Compared to values in **Table 2**, the field-based estimates are 1.6 to 2.4 times greater than the modelled carbon stocks. This difference between model and field data illustrates the extent to which country or local conditions can vary from statically-derived average estimates (e.g., IPCC soil carbon values) and modelled values derived at a global extent, e.g., Hamilton and Friess (in review) and

²⁷ Jardine, S. and J. Siikamäki. 2014. "A global predictive model of carbon in mangrove soils." *Environmental Research Letters*, 9:104013.

²⁸ Hiraishi, Takahiko, et al. 2014. *2013 supplement to the 2006 IPCC guidelines for national greenhouse gas inventories: Wetlands*.

²⁹ Alongi, D. et al. 2015. "Indonesia's blue carbon: a globally significant and vulnerable sink for seagrass and mangrove carbon." *Wetlands Ecology and Management*, 24:3-13.

Table 2. Estimates of mangrove and soil (1 m depth) carbon stocks in 2012.

Country	Hamilton mangrove and soil model (MMt C)	Hutchinson biomass model w/ IPCC SO mangrove, organic (MMt C)	Hutchinson biomass model w/ IPCC SO mangrove, aggregate (MMt C)	Hamilton mangrove and soil model (MMt CO ₂ eq.)	Hutchinson biomass model w/ IPCC SO mangrove, organic (MMt CO ₂ eq.)	Hutchinson biomass model w/ IPCC SO mangrove, aggregate (MMt CO ₂ eq.)
Indonesia	1,253.8 (14.9)	1,900.3	1,646.4	4,597.3 (54.6)	6,967.6	6,036.8
Malaysia	254.2 (3.1)	455.8	395.5	932.0 (11.2)	1,671.2	1,450.0
Philippines	102.3 (0.9)	157.2	135.3	375.2 (3.4)	576.5	496.1
Thailand	90.0 (1.1)	148.7	127.5	330.0 (4.0)	545.4	467.5
Viet Nam	33.0 (0.5)	58.0	49.4	121.0 (2.0)	212.7	181.1
Cambodia	15.1 (0.1)	35.9	30.8	55.4 (0.5)	131.5	112.8
China	5.2 (0.1)	10.7	9.0	18.9 (0.4)	39.2	32.9
Brunei Darussalam	0.3 (0.007)	11.6	10.2	1.3 (0.03)	42.7	37.4
Timor-Leste	0.2 (0.002)	n.d.	n.d.	0.9 (0.006)	n.d.	n.d.
Japan	0.1 (0.007)	n.d.	n.d.	0.4 (0.03)	n.d.	n.d.
Singapore	0.07 (0.008)	n.d.	n.d.	0.2 (0.03)	n.d.	n.d.
Total	1,754	2,778	2,404	6,433	10,187	8,815

Note: Values in parentheses represent standard error; n.d. = no data.

Hutchinson et al. (2013). While model-based values may be sufficient for a regional estimate of carbon stocks, more accurate carbon stocks and carbon stock change are obtained through country-specific data. Addressing this discrepancy is a key consideration for future blue carbon research, highlighting the need to obtain better country-specific stock values, which ultimately will help to better calibrate global models.

To assess how rapidly mangroves are being lost across the region, two approaches are applied. The first approach calculates total percent of mangrove area change between 2000 and 2012 and yearly rates of change based on the Hamilton and Casey (2016) dataset (**Table 3**). Malaysia, Cambodia and Indonesia have the largest percentage loss of mangroves, ranging from 3 to 5%. Singapore, Japan and Brunei Darussalam had the lowest mangrove loss, with less than 1% lost over a 12-year period.

Table 3. Estimates of mangrove loss between 2000 and 2012.

Country	Percent Mangrove Loss			
	MFW ^a		BIOME ^b	
	2000 - 2012	Annually	2000 - 2012	Annually
Malaysia	4.89%	0.41%	13.23%	1.10%
Cambodia	4.48%	0.37%	11.87%	0.99%
Indonesia	3.11%	0.26%	7.68%	0.64%
Thailand	2.44%	0.20%	0.75%	0.06%
China	2.11%	0.18%	2.49%	0.21%
Philippines	1.28%	0.11%	2.10%	0.17%
Timor-Leste	1.28%	0.11%	1.20%	0.10%
Viet Nam	1.15%	0.10%	3.37%	0.28%
Brunei Darussalam	0.79%	0.07%	7.68%	0.64%
Japan	0.76%	0.06%	1.94%	0.16%
Singapore	0.00%	0.00%	2.46%	0.21%

^a Mangrove Forests of the World dataset; ^b Terrestrial Ecosystems of the World dataset (Hamilton and Casey 2016)

A second approach developed by Richards and Friess is also applied.³⁰ The analysis incorporates global forest change data from 2000 to 2012 combined with analysis of satellite imagery to describe the extent of mangrove deforestation and the type of land uses created with deforestation throughout East Asia. China, Singapore and Japan were not included in their analysis.³¹ As with the prior approach, most mangrove deforestation identified occurred in Malaysia and Indonesia. This loss was primarily due to oil palm plantations and aquaculture, respectively. Across all countries examined,

aquaculture was the primary driver of mangrove deforestation, followed by oil palm plantations (Table 4).

In summary, East Asia has some of the largest coverage and biomass of mangroves globally due to the location and climate of its countries. Indonesia and Malaysia are two of the top mangrove-holding countries globally, and have some of the greatest deforestation rates as such, primarily for aquaculture but also palm oil where mangroves meet terrestrial peatlands.

Table 4. Type and amount of mangrove change (in hectares) between 2000 and 2012.⁹

Country	Type and amount of change (ha)							
	Aquaculture	Recent deforestation	Oil palm	Rice	Erosion	Urban	New Growth Mangrove	New Terrestrial Forest
Indonesia	29,591	3,123	9,578	45	2,433	1,127	13,768	1,241
Malaysia	2,764	1,272	7,193	15	163	2,412	3,316	1,701
Thailand	379	151	1,403	195	1	503	179	693
Philippines	523	465	158	12	12	38	105	111
Cambodia	338	457	108	18	1	56	119	122
Viet Nam	112	8	3	55	17	332	3	1
Brunei Darussalam	14	7	13	0	0	8	6	0
Timor-Leste	0	0	0	1	2	0	0	0

Note: no data are available for China, Singapore, and Japan.

Tidal Marsh Ecosystems

Tidal marshes are found at latitudes that are too cold for mangroves to tolerate, typically latitudes greater than 30°N and 30°S. Therefore, in East Asia, tidal marshes occur primarily in China, Japan, DPR Korea and RO Korea. The term “tidal marsh” includes salt, brackish and freshwater marshes, though most tidal marsh research has occurred in salt marshes and thus is the focus of this study.

Tidal marsh ecosystems across East Asia are far less well mapped and studied than mangroves, and few academic papers or reports have been published regarding their distribution and carbon stocks. The technologies exist to map tidal marshes but, to date, no comprehensive datasets have been assembled within the region.³² This study was unable to find any documentation on tidal marsh distribution, carbon stocks or losses for Japan, DPR Korea and RO Korea. This is a significant information gap, yet capacity exists to fill the gap.

³⁰ Richards, D. & D. Friess. 2016. “Rates and drivers of mangrove deforestation in Southeast Asia, 2000-2012.” *PNAS*, 113:344-349.

³¹ Hansen, M. et al. 2013. “High-resolution global maps of 21st century forest cover change.” *Science*, 342:850-853.

³² Zuo, P. et al. 2012. “Distribution of *Spartina* spp. along China’s coast.” *Ecological Engineering*, 40:160-166.


The UNEP World Conservation Monitoring Center (WCMC) compiled the first global dataset specifically on salt marsh distributions; however, the data remain limited in terms of extent, especially within tropical areas where mangroves dominate.³³ The Nature Conservancy and WCMC estimated the relative abundance of salt marshes based on the proportion of coastline within a given region and created an abundance index (**Figure 2**).³⁴ The data highlight low to moderate salt marsh abundance along coastal China, DPR Korea, RO Korea and Japan. It is possible that mangroves are being classified as salt marshes within Viet Nam, Cambodia and Thailand since mangroves are the dominant vegetation type below 30°N.

Most tidal marshes should occur in China due to its latitudinal range and extent of coastline. Notably, Zuo et al³⁵

describe that 57,344 ha of salt marsh occurred in China as of 1995, half along the Bohai Sea and a third along the Yellow Sea coasts and Changjiang delta. In China, *Spartina anglica*, a salt marsh grass, was introduced in the 1960s as were two others in the 1970s, *Spartina alterniflora* and *Spartina cynosuroides*.⁸ These species out-competed many native species, creating monospecific stands that extend lower into the tidal zone, continuing to significantly change the character of Chinese salt marshes and considered to be invasive by many. Recent attention has been paid to mapping the extent and rate of this change.

Due to lack of data in multi-date estimates of tidal marsh distributions in East Asia, this study is unable to include a comprehensive analysis of tidal marsh gain or loss and the subsequent changes in carbon.

Figure 2. Relative abundance of salt marsh habitat based on proportion of coastline within a region.



³³ Hoekstra, J. M. et al. 2010. The atlas of global conservation: changes, challenges, and opportunities to make a difference. Ed. J. L. Molnar. Berkeley: University of California Press.

³⁴ *Ibid.*

³⁵ Shi-Jun, Y. and C. Ji-yu. 1995. "Coastal salt marshes and mangrove swamps in China.


In the absence of data, it is speculated that no more than 60,000 ha of tidal marsh remains across the region. There is a critical need for a tidal marsh change detection analysis to occur, similar to the mangrove analyses in the previous section, spanning at least a ten-year period. However, specifically in China, it was reported that 708,200 ha of salt marshes have been drained, diked or otherwise modified for aquaculture, rice cultivation and development.³⁶ While data are not available for tidal marsh plant carbon stocks, applying IPCC default values for soil organic carbon to tidal marshes with a mixture of organic and mineral soil,³⁷

total soil carbon emissions are estimated to be 180.6 MMt C (or 662.2 MMt CO₂).

Seagrass Ecosystems

Seagrass beds in the South China and Java Seas are some of the most diverse globally (**Figure 3**).^{38,39} Throughout the seas of East Asia, 19 species of seagrass have been identified, or roughly a third of the total number of seagrass species worldwide. But, of all blue carbon ecosystems, seagrass beds are the least documented and mapped within East Asia. This represents a major information gap.

Figure 3. Number of seagrass species located in coastal waters of East Asia.



³⁶ *Ibid.*

³⁷ 255 t C ha⁻¹; (95% CI: 254, 297); IPCC 2014

³⁸ Hoekstra, J. M. et al. 2010. The atlas of global conservation: Changes, challenges, and opportunities to make a difference.

³⁹ Hutomo, M. and M. Moosa. 2005. "Indonesian marine and coastal biodiversity: present status." *Indian Journal of Marine Sciences*, 34:88-97.

Seagrass beds provide food and habitat for many species including sea turtles, dugongs and sea-horses and many commercially important fish species, and they are some of the most threatened habitats due to their close proximity to the shore. Not only are they affected by natural events, such as typhoons and other storms, they are also heavily impacted by suspended sediment, eutrophication, heavy metals, aquaculture and fishing and trawling.

Seagrass beds are also the most difficult blue carbon ecosystem to determine distributions for due to their subtidal location (i.e., they are underwater). Remote sensing techniques commonly used for mapping terrestrial and wetland habitats generally do not work for seagrasses due to the constant presence of water, which inhibits use of most satellite or airborne sensors. Therefore, most seagrass surveys are conducted through field observation, often using SCUBA diving and manual delineation with GPS units.

Some of the largest areas of seagrass beds located within the South China Sea occur near Cape Bolinao, Philippines (22,400 ha); Kampot Province, Cambodia (25,200 ha); Phu Quoc, Viet Nam (12,500 ha); and East Bintan, Indonesia (2000 ha).⁴⁰ Along coastal southern China, seagrass beds totalling 2,400 ha have been surveyed: 910 ha, 690 ha and 871 ha in Guangdong, Guangxi and Hainan provinces, respectively, with very small patches in Hong Kong.⁴¹ Approximately 78 seagrass beds have been identified in Malaysia,⁴² with 316 ha identified in peninsular Malaysia.⁴³ Thailand and Viet Nam have at least 6,850 and 44,000 ha of known sea-

grass beds. In the Philippines, a country-wide survey is currently underway (M. Fortes, personal communication). No nationwide studies on seagrass area in RO Korea have been conducted, but estimates of eelgrass coverage are between 5,500 and 7,000 ha along the southern coast. In Japan, eelgrass bed coverage was estimated to be 49,600 ha. Estimates of 3,000,000 ha of seagrass beds occur throughout Indonesia, although these have not been thoroughly mapped.⁴⁴ No area estimates are known for DPR Korea, Timor-Leste or Brunei Darussalam. More field-based surveys are needed, along with research into newly-developed remote sensing techniques that could aid in more accurately and extensively identifying seagrass beds.

WCMC compiled a global database of seagrass locations based on a combination of points and mapped boundaries from literature and expert knowledge (**Figure 4**).^{45,46} As noted above, spatial data are lacking for East Asian countries. Notably, there are no data for Brunei Darussalam. However, this might be explained by coastal conditions, such as soil substrate and exposure, which could inhibit growth.⁴⁷

As with tidal marshes, the lack of data on seagrass distributions, both currently and historically, provide challenges to documenting changing carbon storage and emissions. Alongi et al. (2015) conducted a carbon stock assessment of Indonesian seagrasses and estimated that the approximately 3,000,000 ha of seagrass beds contain 368.5 MMt CO₂ within the plants and soil. There is an urgent need for establishing current coverage as a baseline for future change analysis.

⁴⁰ Vo, S. et al. 2013. "Status and trends in coastal habitats of the South China Sea." *Ocean & Coastal Management*, 85:153-163.

⁴¹ Huang, X. et al. 2006. "Main seagrass beds and threats to their habitats in the coastal sea of South China." *Chinese Science Bulletin*, 51 Supp. II 136-142.

⁴² Bujang, J. et al. 2006. "Distribution and significance of seagrass ecosystems in Malaysia." *Aquatic Ecosystem Health & Management*, 9:1-14.


⁴³ Green, E. and F. Short. 2003. *World atlas of seagrasses*. Prepared by the UIMEP World Conservation Monitoring Centre. University of California Press, Berkeley, USA.

⁴⁴ Alongi, D. et al. 2015. "Indonesia's blue carbon: a globally significant and vulnerable sink for seagrass and mangrove carbon."

⁴⁵ Short, F. et al. 2007. "Global seagrass distribution and diversity: a bioregional model." *Journal of Experimental Marine Biology & Ecology*, 350.1:3-20.

⁴⁶ UNEP-WCMC, Short FT. 2016. *Global distribution of seagrasses (version 4.0). Fourth update to the data layer used in Green and Short (2003)*. Cambridge (UK): UNEP World Conservation Monitoring Centre. <http://data.unepwcmc.org/datasets/7>. Accessed 2016 November 17.

⁴⁷ Fortes, M. 1990. *Seagrasses: a resource unknown in the ASEAN region*. Vol. 5. WorldFish.

Figure 4. Seagrass beds that have been mapped in East Asia.

as intertidal or subtidal habitat can be derived. Much of this area will once have been tidal wetlands historically, now converted through human actions to other lands uses, such as agriculture or settlement. Total area by country is given in **Table 5** and country-specific maps are available in the Appendix. Notably, China contains the largest extent of coastal floodplain.

At 3,359,041 ha, the total area of converted and diked former tidal wetlands across East Asia is substantial (by comparison, the drained coastal wetland area of the conterminous United States is 529,551 ha).⁵⁰

Large rivers flowing from the Himalayas carry sediment, building vast deltas and coastal floodplains, notably at the mouths of rivers Yellow, Yangtze, Pearl, Red and Mekong. Expansive coastal plains also formed at the mouth of smaller rivers, such as the Chao Phraya, and along the island coastal margins of Indonesia. These low-lying flat lands are both highly attractive for human occupation, but they are also highly vulnerable to, and costly to protect from, flooding.

Over time, vulnerability to flooding will increase in delta areas. This is due to a combination diked areas being isolated from river sediment supply that once compensated for natural high rates of subsidence that occur across delta areas, coupled with rising water levels as sea level rises and river flood flows increase due to climate change.^{51,52}

Historic Loss of Blue Carbon Ecosystems

While distributions over time are not mapped for most blue carbon ecosystems in East Asia, long-term historic loss of mangroves and tidal marshes can be estimated.⁴⁸ Using satellite-derived elevation data,⁴⁹ the area of land now at or below mean sea level that was not already classified

increase in delta areas. This is due to a combination diked areas being isolated from river sediment supply that once compensated for natural high rates of subsidence that occur across delta areas, coupled with rising water levels as sea level rises and river flood flows increase due to climate change.^{51,52}

⁴⁸ The calculated area is a reasonable approximation of former tidal wetland area. It will include some areas of former unvegetated tidal and subtidal wetlands that could accentuate the estimate, but also filled tidal wetland areas now above high tides that would detract from the calculated area.

⁴⁹ Shuttle Radar Topography Mission (SRTM) with 90m resolution and vertical accuracy of 1m. SRTM-derived water bodies and mangrove areas (per Giri et al. 2011) were masked from the dataset so as not to overestimate the floodplain level area.

⁵⁰ Crooks, S. et al. 2016. *Bringing Coastal Wetlands into the U.S. Inventory of Greenhouse Gas Emissions and Sinks*. Presentation at Restore America's Estuaries Summit, New Orleans December 11, 2016.

⁵¹ Syvitskiet, J. et al. 2009. "Sinking Deltas Due to Human Activities. *Nature Geoscience*, 2:681-686.

⁵² Arnell, N.W. and Gosling, S. N. 2016. "The impact of climate change on river flood risk at the global scale." *Climate Change*, 134(1):387-401.

Table 5. Estimated soil carbon emissions from converted former coastal tidal wetlands.

Country	Land at or below mean sea level (ha)	Estimated soil carbon emissions, top meter of soil (MMt C)	
		mean	95% CI
China	1,813,766	462.5	460.7, 538.7
Viet Nam	849,966	328.1	298.3, 360.4
Indonesia	301,238	116.3	105.7, 127.7
Japan	233,694	59.6	59.4, 69.4
Philippines	68,194	26.3	23.9, 28.9
Thailand	53,441	20.6	18.8, 22.7
RO Korea	12,649	3.2	3.2, 3.8
Cambodia	12,056	4.7	4.2, 5.1
DPR Korea	7,368	1.9	1.9, 2.2
Malaysia	5,557	2.1	2.0, 2.4
Singapore	644	0.2	0.2, 0.3
Timor-Leste	464	0.2	0.2, 0.2
Brunei Darussalam	4	0.0017	0.0015, 0.0018
Total	3,359,041	1,026	1022.0, 1195.1

Note: Estimated soil carbon emissions based on IPCC values for aggregated mineral and organic soils; all values are for mangroves except for China, Japan, RO Korea and DPR Korea, which are for tidal marshes.

These former tidal wetland areas are also historic sources of carbon emissions.⁵³ Using these estimates of former mangrove and tidal marsh areas and IPCC approaches for quantifying soil stock change, estimated emissions from conversion of mangroves and tidal marshes are provided for each country. Cambodia, RO Korea, Brunei Darussalam, Timor-Leste, Singapore, DPR Korea and Malaysia all had less than 5 MMt soil carbon lost, while China and Viet Nam had 462 and 328 MMt soil carbon lost, respectively. Historic coastal land-use for aquaculture, agriculture and development are the primary drivers of this change.

These estimates of soil carbon emissions are very conservative in that they assume emissions are limited to the top one meter of soils. Drained organic soils will continue to emit carbon dioxide back to the atmosphere until the resource of carbon is exhausted or the soils are reflooded. This depth can be much greater than one meter. Such

an example of long term emissions can be found in the Sacramento-San Joaquin Delta of California, where soils converted to agricultural land between 1850-1920 now lie over 7 meters below sea level and have emitted around one billion tons of carbon dioxide since conversion.⁵⁴ These emissions in the Sacramento-San Joaquin Delta continue to this day.

There is a need to understand more clearly the distribution of organic soils on converted coastal lands, particularly deltas, how the current land management practices influence emissions and whether modified management practices can reduce emissions without impacting significantly uses. Over time, as sea level rises and lands subside, maintaining existing land uses on delta systems will become increasingly challenging. Delta systems offer high potential for wetland restoration if lands are reconnected to river channels and sediment delivery. Through this restoration process emissions stop.

⁵³ Crooks, S. et al. 2011. "Mitigating climate change through restoration and management of coastal wetlands and near-shore marine ecosystems: challenges and opportunities."

⁵⁴ *Ibid.*



Photo by SCA/Crooks

3

Blue Carbon as a Climate Intervention Concept: UNFCCC, Kyoto Protocol and the Paris Agreement

International climate policy has long been concerned with formulating emission reduction objectives and (to a lesser extent) resilience goals, but it has also provided an ever-increasing number of instruments—some specifically designed for non-government entities and some triggering considerable funding—that help achieve the objectives and goals set.

The importance of mangroves for designing climate change *adaptation* instruments has long been recognized by scientists, policymakers and civil society groups across countries.⁵⁵ It was not until recently, however, that the specific value and role of mangroves, tidal marshes and seagrasses for climate change *mitigation* were addressed by climate policy experts).

Widely regarded as one of its greatest weaknesses, the Kyoto Protocol focused almost exclusively on climate change mitigation, yet without providing for any instruments to combat emissions from deforestation and ecosystem degradation.⁵⁶ Twenty years on, with the Paris Agreement freshly entered into force, the situation is quite different. The constructive synergies and overlaps between blue carbon interventions

and adaptation efforts are beginning to inform policymakers worldwide. Conservation, restoration and sustainable use of sensitive coastal ecosystems are increasingly understood in their multiple roles to support livelihoods, store carbon and increase resilience against climate change. In its preamble, the Paris Agreement notes the “importance of ensuring the integrity of all ecosystems, including oceans, and the protection of biodiversity”, and a full article is reserved for the Parties’ commitment to “conserve and enhance, as appropriate, sinks and reservoirs of greenhouse gases”.

While the majority of countries, in their climate change strategies and their international commitments—intended nationally determined contributions or “INDCs”—made in anticipation of the Paris Agreement, still largely treat coastal conservation and restoration efforts exclusively within the scope of adaptation and resilience, there are early movers and pioneers that have explicitly mentioned coastal blue carbon ecosystems as part of their mitigation efforts.⁵⁷ An increasing number of governments and non-government institutions have also started to conceptualize specific opportunities presented when linking blue

⁵⁵ McLeod, Elizabeth and Rodney V. Salm. 2006. “Managing mangroves for resilience to climate change.” World Conservation Union (IUCN), Gland. http://cmsdata.iucn.org/downloads/managing_mangroves_for_resilience_to_climate_change.pdf. Accessed 2016 December 8.

⁵⁶ Joosten, H. et al. 2016. “Peatlands, forests and the climate architecture: setting incentives through markets and enhanced accounting.” <https://www.umweltbundesamt.de/publikationen/peatlands-forests-the-climate-architecture-setting>. Accessed 2016 December 8.

⁵⁷ For an overview of blue carbon country commitments in INDCs see D. Herr / E. Landis, Coastal blue carbon ecosystems (IUCN / TNC 2016).

carbon interventions with conservation finance⁵⁸ and payment-for-ecosystem services (“PES”) schemes,⁵⁹ on the one hand, and new climate finance tools, such as results-based finance,⁶⁰ blue bonds⁶¹ and debt-swap-for-nature agreements,⁶² on the other hand. Several country-specific efforts show that there is growing interest in including the carbon value of these three ecosystems in national policy, planning and decision-making.⁶³

History I: Between Mitigation and Adaptation

Climate change policymaking has long been split between climate change mitigation—reducing CO₂ and other greenhouse gases (measured by their warming potential as CO₂ equivalent or CO₂ eq)—and climate change adaptation. This split was never simply a breakdown of tasks. Rather, it always represented a choice of objectives, either combating climate change or dealing with its consequences, particularly when focusing on the former allowed for a clear allocation of responsibility of action (for industrialized countries). This paradigmatic choice was always a curious one, not based in the United Nations Framework Convention on Climate Change (UNFCCC), which formulated a functional relationship: to stabilize emissions of greenhouse gases (GHG) *in order to allow* ecosystems to adapt naturally (Article 2 UNFCCC). Yet, it drove the negotiations of the Kyoto Protocol and was very effective in ensuring that for 20 years of climate change policy making, the focus was on climate change mitigation and what industrialized countries could do on that front. This led to a prioritization of addressing industrial production

and the generation and use of energy, to the disadvantage of land-use matters—“LULUCF” in the terminology of climate negotiations: land use, land-use change and forestry—at large.

History II: Blue Carbon Accounting Rules and Practice

This said, the focus on mitigation and energy was also a pragmatic one. While it is fairly easy to measure and calculate GHG emissions from industrial installations, measuring and calculating GHG fluxes from ecosystems is far more complex, involving questions of agency, human-induced or natural, as well as temporality (permanence of sequestration). Thus, largely reducing GHG accounting to energy-related activities allowed Parties to the Kyoto Protocol to formulate with certainty historic emissions (using a base year for most countries of 1990) and targets to be reached by 2012.

Still, industrialized countries in calculating GHG emissions had to account for certain land-related activities, such as forestland and cropland management. Following the release of the IPCC Wetland Supplement in 2014 countries were given the opportunity to also recognize emissions and removals with wetland drainage and rewetting. Accounting for these emissions is voluntary, though countries such as the United State and Australia are amongst a few early adopters in applying the Wetland Supplement guidance.

Developing countries, which host the clear majority of blue carbon ecosystems, have no accounting obligations. They are still requested to report GHG emissions

⁵⁸ Huwyler, F. et al. 2014. *Conservation finance: moving beyond donor funding toward an investor-driven approach*. Credit Suisse, WWF, McKinsey & Company: Zurich, Switzerland.

⁵⁹ Locatelli, T. et al. 2014. “Turning the tide: how blue carbon and payments for ecosystem services (PES) might help save mangrove forests.” DOI 10.1007/s13280-014-0530-y. Accessed 2016 December 9.

⁶⁰ For a project-based perspective see S. Crooks et al. 2014. Guiding principles for delivering coastal wetland carbon projects, at http://www.unep.org/pdf/Guiding_principles_for_delivering_coastal_wetland_projects.pdf.

⁶¹ For the concept on “green bonds” see World Bank Treasury at http://treasury.worldbank.org/cmd/pdf/What_are_Green_Bonds.pdf; the issuance of “blue bonds” is under consideration by, among others, the Seychelles, cf. Bloomberg News of 24 January 2016: “Seychelles Plans Blue Bonds to Develop Sustainable Fisheries”, at <http://www.bloomberg.com/news/articles/2016-01-24/seychelles-plans-blue-bond-sale-to-develop-sustainable-fisheries>.

⁶² Resor, J. 2010. “Debt-for-nature swaps: a decade of experience and new directions for the future.” <http://www.fao.org/docrep/w3247e/w3247e06.htm>. Accessed 2016 December 9.

⁶³ Cf. the “Sustainable Landscapes in Eastern Madagascar” Project, a cooperation between the European Investment Bank and Conservation International, with initial funding provided by the Green Climate Fund to prepare green / blue bond issuance; see also UNDP’s project for Viet Nam (“Improving the resilience of vulnerable coastal communities to climate change related impacts in Viet Nam”), which includes mitigation and adaptation activities rooted in mangrove rehabilitation (funded by the Green Climate Fund).

as part of their national inventories and, since recently, in the form of biennial update reports (BURs). But reporting standards remain much weaker than the target-oriented accounting standards. While the methodological reporting approach is improving, the overall quality of data remains paltry. Mangrove-related data is nominally included in those countries that consider mangroves as forest,⁶⁴ but the calculation for most areas is incomplete. The IPCC recently issued a dedicated section on accounting for emissions from wetlands, including mangrove forests, tidal marshes and seagrass meadows in 2013,⁶⁵ and these up-to-date accounting guidelines are only now starting to be applied by pioneering countries. Blue carbon ecosystems other than mangrove forests are usually not covered.

CDM: Mangrove Afforestation and Reforestation Approaches

The Clean Development Mechanism (CDM) is the most prominent of the so-called “flexible mechanisms” of the Kyoto Protocol. It allows industrialized countries to invest in emission reduction or sequestration projects in developing countries in exchange for the issuance of credits, or Certified Emission Reductions (CERs), which can be used by industrialized countries to offset their emissions and to comply with the Kyoto target.

The CDM is highly restrictive when it comes to land-use based projects. It permits afforestation and reforestation projects only (excluding conservation projects proper) and it issues temporary credits only in acknowledgement of the risk that a forest, once planted, may burn, be chopped down or otherwise degrade (Joosten et al. 2016). Still, the CDM led to the development of more than

10 ecosystem-based accounting methodologies (for example on “Afforestation and reforestation of degraded mangroves”)⁶⁶ and some 50 projects worldwide, among them a “Small-scale and low-income community-based mangrove afforestation project on tidal flats” in Riau Island Province, Indonesia.⁶⁷

Voluntary Carbon Markets

Voluntary carbon standards spread where the CDM did not provide action formats, notably in the area of ecosystem conservation. The Verified Carbon Standard (VCS), in particular, consolidated blue carbon-related project approaches. Under its sectoral scope “Wetlands”, it offers a methodology for coastal wetland creation as well as a methodology for tidal wetland and seagrass restoration.⁶⁸

While the overall value of the voluntary carbon markets has been stable for some years, additional demand is expected to come from the aviation industry, which in 2016 paved the way for a carbon offsetting instrument. In the framework of the International Civil Aviation Organization (ICAO), countries committed to freeze aviation emissions from 2020 onwards.⁶⁹ To compensate for increasing emissions, a dedicated mechanism will be set up: the Carbon Offsetting and Reduction Scheme for International Aviation (“CORSIA”). Some 65 countries representing more than 80% of international air traffic have signed up so far and countries and airlines are given discretion when preparing for the piloting phase (starting, formally, in 2020). The Verified Carbon Standard, in particular, hopes for clearance of VCS projects under CORSIA.

⁶⁴ The two most commonly used definitions of forest are those used by the Food and Agriculture Organization of the United Nations (FAO) – forest means all land spanning more than 0.5 hectares with trees higher than 5 meters and a canopy cover of more than 10% – and the Kyoto Protocol Marrakesh Accords, UNFCCC (2001) COP.7: forest requires minimum area between 0,05 to 1 hectare, a minimum height at maturity of between 2-5 meters, and a crown cover of between 10-30%; countries may freely decide within these scopes.

⁶⁵ Hiraishi, Takahiko, et al. 2014. *2013 supplement to the 2006 IPCC guidelines for national greenhouse gas inventories: Wetlands*.

⁶⁶ UNFCCC. [n.d.]. “AR-AM0014: Afforestation and reforestation of degraded mangrove habitats --- Version 3.0.” <https://cdm.unfccc.int/methodologies/DB/KMH608T6RL3P5XKNBQE2N359QG7KOE>. Accessed 2016 December 12.

⁶⁷ UNFCCC. [n.d.]. “Small-scale and low-income community-based mangrove afforestation project on tidal flats of three small islands around Batam City, Riau Islands Province, Republic of Indonesia.” <https://cdm.unfccc.int/Projects/Validation/DB/Q94TKNG9V6GRFOTRKLWDV2GHBCFWW/view.html>. Accessed 2016 December 12.

⁶⁸ See the VCS Methodology for Tidal Wetland and Seagrass Restoration, <http://database.v-c-s.org/methodologies/methodology-tidal-wetland-and-seagrass-restoration-v10>.

⁶⁹ ICAO. 2016. “Draft text for the report on agenda item 22 (Section on Global Market-based Measure Scheme).” http://www.icao.int/Meetings/a39/Documents/WP/wp_462_en.pdf. Accessed 2016 December 12.

REDD: Overlapping Concepts

Over recent years, discussions on how to reduce emissions from deforestation and forest degradation (REDD+) have emerged as one of the main negotiation blocks between UNFCCC Parties.⁷⁰ A comprehensive framework – the Warsaw Framework for REDD+⁷¹ – was established, and the Paris Agreement specifically encourages Parties to take actions to promote REDD+.⁷² While REDD+ remains a complex instrument, important progress has been made over the years on many fronts and many technical details, including the calculation of forest emission reference levels, REDD+ safeguards, REDD+ implementation phases and jurisdictional (sub-national) approaches. A range of bilateral and multilateral activities have supported REDD+ development, among them the *United Nations Collaborative Initiative on Reducing Emissions from Deforestation and forest Degradation* (UN REDD) and the World-Bank-managed *Forest Carbon Partnership Facility* (FCPF). Some fifty, mostly tropical, countries—including the majority of PEMSEA partner countries—have started building country-wide REDD/REDD+ implementation frameworks, with financial support from developed countries. By 2015, almost \$9 billion USD had been pledged by donors to support REDD+ activities.⁷³ For an overview of funding facilities with a special focus on REDD+ see the later discussion on linking climate finance and blue carbon (Section IV).

REDD+ and blue carbon interventions share a number of characteristics. They have habitat overlap, as far as mangroves are concerned (and as long as countries consider mangroves as forest, which most, but not all, do). They share many direct drivers of degradation, notably agriculture (for blue carbon habitats, aquaculture, in particular), timber extraction and development (construction), as well as indirect causes, including land tenure related challenges.⁷⁴ Governments increasingly design REDD+ projects and toolkits for coastal environments, sometimes highlighting the importance of mangrove forests.⁷⁵ Nevertheless, REDD+ remains a slow process, and most countries have not yet tapped into the scheme's full potential.⁷⁶

It is also noted that REDD+ engagement often holds a certain level of ambiguity for blue carbon ecosystems, which in turn can impede project-level action. First, it is often not clear to what extent blue carbon ecosystems are covered by a country's REDD+ policy framework. At first sight, several countries, including Indonesia, Malaysia and Viet Nam, cover mangroves for the purpose of defining their REDD+ reference levels. Yet, looking deeper, these countries mostly include above-ground biomass only, leaving the enormous below-ground carbon sink in limbo.⁷⁷ Then, second, as the primary focus of REDD+ is nation-wide or jurisdictional engagement, the role of REDD+ projects remains vague, and carbon project developers have the (at

⁷⁰ La Viña, A. et al. 2016. "History and future of REDD+ in the UNFCCC: issues and challenges." In *Research Handbook on REDD+ and International Law*. Edited by C. Voigt. <http://www.e-elgar.com/shop/eep/preview/book/isbn/9781783478316/>. Accessed 2017 January 25. See further on REDD and blue carbon and blue carbon commonalities Herr et al. 2017. *Pathways for implementation of blue carbon initiatives*, Aquatic Conservation (forthcoming).

⁷¹ Decisions 9-15 of Warsaw Framework for REDD-plus (COP 19, Warsaw), http://unfccc.int/land_use_and_climate_change/redd/items/8180.php. Accessed 2016 December 12.

⁷² Article 5.2 of the Paris Agreement.

⁷³ Lee, D. and T. Pistorius. 2015. "The Impacts of International REDD Finance." http://www.climateandlandusealliance.org/wp-content/uploads/2015/09/Impacts_of_International_REDD_Finance_Report_FINAL. Accessed 2016 December 15.

⁷⁴ Rotich, B. et al. 2016. *Where land meets the sea. A global review of the governance and tenure dimensions of coastal mangrove forests*.

⁷⁵ For a case study in Kenya see L. King, *Notes From The Field: Including mangrove forests in REDD+* (CDKN 2012), at <http://cdkn.org/wp-content/uploads/2012/12/Notes-from-the-field-Lesley-King-1.pdf>; for a case study in Colombia, see <http://bioredd.org/projects/>; for a feasibility study on REDD+ and the Japanese Joint Crediting Mechanism (JCM) see Japanese Ministry of Economy, Trade and Industry. 2015. *Mangrove REDD+ Reduction of CO2 Emission and Enhancement of Carbon Fixation*, at http://www.meti.go.jp/meti_lib/report/2015fy/000139.pdf.

⁷⁶ For a case study on different countries with stronger and weaker results see Lee/Pistorius, op. cit.

⁷⁷ Malaysia states in its reference level submission that it includes both above-ground and below-ground biomass; however, the assessment team notes that emissions levels of soil organic carbon, among others, may be an "area for future technical improvement" and points to the development of a soil organic database under way in Malaysia, cf. UNFCCC: Report on the technical assessment of the proposed forest reference level of Malaysia submitted in 2014 (17 December 2015), para. 26.

times arduous) responsibility of clarifying whether the REDD+ architecture permits bottom-up development projects in the first place. This said, various countries have started setting up structures for defining the scope of REDD+ projects and the mechanics of their “nesting” is also established under voluntary standards.⁷⁸ The efforts will simplify interventions for blue carbon developers, too.

Blue Carbon and NAMAs

National Appropriate Mitigation Actions (NAMAs) were first introduced under the Bali Action Plan as a vehicle for parties in developing countries to develop nationally specific mitigation actions. NAMAs have since evolved to support the implementation of the nationally determined country contributions (NDCs) in the framework of the Paris Agreement, as well as the 2030 Agenda for Sustainable Development (the UN SDGs) alongside national development strategies (see section IV). The land-use sector offers opportunities for NAMA engagement.⁷⁹

NAMAs are climate change mitigation measures voluntarily proposed by developing country governments to reduce emissions below business-as-usual levels and to contribute to domestic sustainable development. While the regulatory framework remains loose, NAMAs can take the form of regulations, standards, programs, policies or financial incentives.

Key categories any NAMA must address relate to governance (central coordination), mitigation output (measurable reduction of CO₂eq. and/or CO₂eq.

sequestration), coherence and impact within the country policy framework and its sustainability goals, planning and timing, as well as support (technological, capacity, financial, etc.) sought from domestic and/or international sources.

Donors also often point to the need for any NAMA to go beyond project-based approaches both in terms of quantity (mitigation output) and quality (underlying ambition of the host country), to help transform the country towards low-carbon development (trajectory impact), and to work towards long-term sustainable economies. The integration of public and private sector institutions is seen in this regard as a priority.⁸⁰

While the main focus so far has been on NAMA scoping exercises and preparatory assessments, and while key donors, including the Green Climate Fund, are yet to establish a NAMA funding portfolio, a few initiatives and facilities, chiefly the “NAMA Facility”, founded by Germany and the UK and increasingly attracting other donors, have started implementing, and delivering on, NAMAs.

More than 170 NAMAs are under preparation or implementation worldwide. Some 16 NAMAs have received a funding commitment for full implementation.⁸¹ Asia has been slower than other regions to embrace the NAMA concept, but has recently caught up. NAMAs under development in the land-use sector of PEMSEA partner countries include the “NAMA for the Sustainable Development of Peatland in Indonesia” (Indonesia);⁸² “Adaptation and Mitigation Activities in Philippine Rice Cultivation” (Philippines);⁸³ and a

⁷⁸ Pearson, T. et al. 2016. “Guidance Document: Options for Nesting REDD+ Projects.” (Fundación Natura Colombia 2016).

http://www.v-c-s.org/wp-content/uploads/2016/07/Nesting-Options-1-Jul_Eng_final.pdf. Accessed 2016 December 18.

⁷⁹ FAO. 2015. “Learning Tool on Nationally Appropriate Mitigation Actions (NAMAs) in the agriculture, forestry and other land use (AFOLU) sector.” <http://www.fao.org/3/a-i4642e.pdf>. Accessed 2016 December 20.

⁸⁰ Bosquet, Michelle et al. 2016. “Status Report on Nationally Appropriate Mitigation Actions (NAMAs): Mid-Year Update Report 2016.” Edited by Charlotte Cuntz et al. <https://www.international-climate-initiative.com/fileadmin/Dokumente/2016/NAMA-Status-Report-June-2016.pdf>. Accessed 2016 December 20.

⁸¹ *Ibid.*

⁸² FAO (2015), op. cit.

⁸³ NAMA. 2016. “Adaptation and mitigation initiatives in Philippine rice cultivation.” <http://naneanews.org/news/2015/03/05/climate-change-adaptation-oriented-nama-option-for-the-rice-sector-in-the-philippines/>. Accessed 2017 January 25.

“NAMA on Sustainable Charcoal Value-Chains in Cambodia” (Cambodia).⁸⁴ Viet Nam is currently setting up a NAMA framework with a sub-window for LULUCF activities.⁸⁵ In other parts of the world, the first blue carbon NAMAs are under development. The Dominican Republic has registered a NAMA on mangrove conservation and restoration (implementa-

tion ongoing).⁸⁶ Ecuador is considering a “Mangrove Blue Services NAMA” that would combine the development of a science and knowledge infrastructure, the establishment of a programmatic roundtable in public-private partnership (inspired by, among others, the Roundtable for Palm Oil) and a community-based management scheme.

⁸⁴ Association GERES. 2017. “NAMA on sustainable charcoal value-chains in Cambodia.” <http://www.geres.eu/en/our-actions/item/481-nama-on-sustainable-charcoal-value-chains-in-cambodia>. <http://www.geres.eu/en/our-actions/item/481-nama-on-sustainable-charcoal-value-chains-in-cambodia>. Accessed 2017 January 25.

⁸⁵ Nguyen Van Huy. [n.d.]. “Towards NAMA/MRV Readiness in Vietnam.” [presentation]. https://unfccc.int/files/focus/mitigation/application/pdf/towards_nama_readiness_invietnam.pdf. Accessed 2017 January 25.

⁸⁶ Blue Carbon NAMA: Conserve and Restore Mangroves in the Dominican Republic, registered with the UNFCCC as a NAMA seeking international support, at [22](http://www4.unfccc.int/sites/nama/_layouts/un/fccc/nama>NamaSeekingSupportForPreparation.aspx?ID=115&viewOnly=1.</p></div><div data-bbox=)



Photo by SCA/Crooks

4

Implementing Blue Carbon Under Nationally Determined Contributions (NDCs)

The Paris Agreement rests on a bottom-up architecture, under which each country is bound to implement mitigation (and arguably adaptation) actions to the extent only that they have been formulated by the country itself and included in the country's so-called Nationally Determined Contribution (NDCs). Prior to the twenty-first Conference of the Parties at Paris in late 2015 ("COP 21"), Parties had submitted what was then, in anticipation of the Paris Agreement, referred to as "intended" NDCs ("INDCs"). A country's INDC becomes its first formal NDC upon ratification of the Paris Agreement by that country (unless it explicitly decides otherwise).⁸⁷ The UNFCCC Secretariat acts as registrar of all NDCs.⁸⁸ At the time of writing, all PEMSEA partner countries except for Brunei Darussalam, Cambodia, the Philippines and Timor-Leste had ratified the Paris Agreement. None had opted out of the INDC transfer mechanism; thus, at the time of writing, those countries that had ratified had formal NDCs in place.

NDC Commitments by Countries in East Asia

The INDC/NDC process has proven enormously successful as a framework for inclusive action and in gathering sup-

port across countries defying the old divide between developed and developing. However, there has been little formal guidance for countries in preparing INDCs/NDCs, and they differ substantially in scope, methodological approach, focus, type of target and ambition.⁸⁹ Only one country from the PEMSEA country bloc, Japan, has formulated a cross-sectoral (100% of activities, including LULUCF), absolute target, in line with other industrialized countries. Japan's target itself (26% of reduction compared with its emissions levels in 2013) has come under critique, however, as "inadequate" in ambition.⁹⁰ Other countries, notably China, Malaysia and partially Viet Nam, have formulated reduction targets related to the amount of CO₂ emitted per unit of GDP. It is noted, in this context, that the role of LULUCF is not yet entirely clear. Malaysia integrates LULUCF emissions in its baseline emissions. Viet Nam seems to account at least net sequestration gains to its target. China's NDC is silent on the matter. Most countries, including most PEMSEA partner countries, set their targets in relation to business-as-usual (BAU) projections, with some countries also addressing LULUCF—or in

⁸⁷ Decision No 1 of the Conference of the Parties (21st session), Dec 1/CP.21, para. 22.

⁸⁸ The registrar is currently an interim solution only, cf. <http://www4.unfccc.int/ndcregistry/Pages/Home.aspx>.

⁸⁹ For an analytical global overview with detailed information on several PEMSEA countries ICF/USAID, Analysis of Intended Nationally Determined Contributions (INDCs), June 2016, at https://www.climatelinks.org/sites/default/files/asset/document/INDC%20White%20Paper%20-%20June%202016_public_RALI.pdf.

⁹⁰ Climate Action Tracker, an independent platform bringing together consultancy firms and experts, namely Climate Analytics, Ecofys, NewClimate Institute and the Potsdam Institute for Climate Impact Research, funded by ClimateWorks, at <http://climateactiontracker.org/about.html>.

the case of Indonesia “LUCF”⁹¹—and others (such as Thailand) remain undecided. All PEMSEA partner countries (except for Japan)⁹² put an emphasis not just on mitigation, but on adaptation. For details, see **Table 6**. In terms of ambition, the (I)NDCs of China, Indonesia and the Philippines have been found “medium” in ambition; Singapore’s INDC received

the same marks as Japan (inadequate). Others have not yet been reviewed following said methodology.⁹³

Almost all PEMSEA partner countries refer, in one way or the other, to blue carbon ecosystems. This reflects the global trend to have blue carbon interventions addressed at the level of (I)NDCs.⁹⁴

Table 6. (I)NDC approaches of PEMSEA partner countries.

Country	INDC/NDC Commitment	LULUCF Accounting	Policies, Measures & Mechanisms (LULUCF)
Brunei Darussalam	<ul style="list-style-type: none"> 63% reduction in total energy consumption by 2035 Reduce fossil fuel demand for inland energy use through a revised power tariff system 	<ul style="list-style-type: none"> LULUCF acknowledged for its sink function (set at 2.63 m CO₂eq. for the year 2010, methodology not referenced) 	<ul style="list-style-type: none"> Allocation of 58% of land area to the “Heart of Borneo” forest conservation area It is aimed to “prevent the possible deterioration of the country’s natural ecosystems” and to “preserve the country’s biodiversity” Transnational partnerships (Malaysia and Indonesia) and work with NGOs (including WWF)
Cambodia	<ul style="list-style-type: none"> 27% reduction in sectors outside LULUCF by 2030, compared with business-as-usual (BAU) Increase forest land cover to 60% of total area by 2030 Both targets are conditional on support 	<ul style="list-style-type: none"> LULUCF considered a net sink In 2010, sequestration value stood at about 18.5 MMt CO₂eq, projected to lower to 7.9 MMt CO₂eq. if no support in the sector is received “Emission reduction” factor assumed per hectare of forest land is 4.7 MMt CO₂eq/ha/year Methodological approach: IPCC Good Guidance Practice INDC acknowledges that “coastal zone resources already face several pressures, including from over-fishing, over-exploitation of forest resources and mangrove ecosystems leading to increased erosion. Climate change adds to these existing challenges through sea level rise, shrinking arable land and decreasing availability of drinking water” 	

⁹¹ “Land use and forestry”. The term apparently seems to exclude status-quo (ongoing) emissions not related to new changes; this may be related to the vast amount of status-quo (ongoing) emissions from drained peatlands.

⁹² *In its NDC, Japan announces the development of a “Plan for Global Warming Countermeasures”, however.*

⁹³ For a recent case study and literature review for several of the PEMSEA countries (albeit without focus on LULUCF) see J. Amponin & J. Evans. 2016. Assessing the Intended Nationally Determined Contributions of ADB Developing Members, ADB Sustainable Development Working Papers Series No 44.

⁹⁴ Herr, D. and E. Landis. 2016. “Coastal blue carbon ecosystems. Opportunities for Nationally Determined Contributions.” https://thought-leadership-production.s3.amazonaws.com/2016/11/12/10/26/14/0d574e9d-3eac-4591-9b8b-57001631b935/BC%20NDCs_FINAL.pdf. Accessed 2017 January 25.

Table 6. (I)NDC approaches of PEMSEA partner countries (cont.)

Country	INDC/NDC Commitment	LULUCF Accounting	Policies, Measures & Mechanisms (LULUCF)
PR China	<ul style="list-style-type: none"> • Peak in CO₂ emissions “around 2030” • 60-65% reduction in CO₂ emissions per unit of GDP by 2030 (compared to 2005 levels) • “Around 20%” increase of share in non-fossil fuels in primary energy consumption (by 2030) • Increase the forest stock volume by 4.5 billion cubic meters from 2005 levels (by 2030) 	<ul style="list-style-type: none"> • While the methodological approach to LULUCF (including its impact on its energy intensity target) remains vague, China stresses the sink effect of its afforestation/reforestation (A/R) targets • LULUCF is mostly acknowledged in its sink capacity, but reference is made to the need to reduce “deforestation-related emissions” 	<ul style="list-style-type: none"> • China aims to “vigorously enhance afforestation, promoting voluntary tree-planting by all citizens... protecting natural forests...” • China further aims to “strengthen the protection and restoration of wetlands and to increase carbon storage capacity of wetlands” • In terms of instruments, China wishes to build on the country’s carbon emission trading pilots “so as to make the market play the decisive role in resource allocation”, and to develop “mechanisms for the reporting, verifying and certificating of carbon emissions” • NDC notes that China “will continue to adapt to climate change by enhancing mechanisms and capacities to effectively defend against climate change risks in key areas such as agriculture, forestry and water resources as well as in cities, coastal and <i>ecologically vulnerable areas</i>.” • China will “enhance resistance to marine disasters and management of coastal zones and improve the resilience of coastal areas against climatic disasters”
DPR Korea	<ul style="list-style-type: none"> • Unconditional 8% reduction of GHGs by 2030 compared with business-as-usual (BAU) • With international support (finance, technology transfer and capacity-building), the reduction could be as much as 40% compared with BAU 	<ul style="list-style-type: none"> • Agriculture, Forestry and Other Land-Use (AFOLU) included in target • Calculation based on 2006 guidelines 	<ul style="list-style-type: none"> • List includes an action target to “manage and develop forest in sustainable manner” by installing tree nurseries, introducing advanced A/R technologies and scaling up technologies for sustainable forest management including agroforestry • Conditional on international support, aim is to replace conventional wood stoves for cooking with efficient wood stoves at rural households; and to scale up agroforestry and sustainable forest management • For the purpose of adaptation, it is envisaged to increase the capacity for integrated management of coastal zones, to construct “infrastructures” such as seawalls and protective facilities in coastal zones, and to “rearrange” population and economic activities • Also, the improvement of ecosystem conservation system(s) in the coastal zone of the Korea West Sea is envisaged

Table 6. (I)NDC approaches of PEMSEA partner countries. (cont.)

Country	INDC/NDC Commitment	LULUCF Accounting	Policies, Measures & Mechanisms (LULUCF)
Indonesia	<ul style="list-style-type: none"> Unconditional 29% reduction by 2030 compared with business-as-usual (BAU) With international support, the reduction target could reach 41% 	<ul style="list-style-type: none"> Target does not include land use as such, but land-use change and forestry (LUCF) NDC recognizes that the country's coastal zone is "affected by the rate of deforestation and forest degradation" and that the "loss of forest ecosystems leads to the loss of critical environmental services, provision of water catchment areas, prevention of erosion and floods" 	<ul style="list-style-type: none"> "REDD+ will be an important component" of the NDC target Indonesia to take an "integrated, landscape-scale approach covering terrestrial, coastal and marine ecosystems" Indonesia will "[protect and restore] key terrestrial, coastal and marine ecosystems"
Japan	<ul style="list-style-type: none"> Reduce absolute sector-wide emissions by 26% by 2030 compared to 2013 emissions Removal target (forests, cropland, grassland, revegetation): 37 MMt CO₂ 	<ul style="list-style-type: none"> LULUCF accounted for: forest-related activities, cropland management, grassland management and revegetation Methodologies "in line with the Guidelines for National [GHG] Inventories prepared by the IPCC, and adopted by the COP" 	<ul style="list-style-type: none"> To promote measures for GHG removals through the "promotion of forest management/forestry industry measures" To promote soil management leading to the increase of carbon stock in cropland To promote revegetation Internationally: 50-100 MMt CO₂ offsets annually to be sourced through the Joint Crediting Mechanism (JCM)
Malaysia	<ul style="list-style-type: none"> 45% reduction of emissions intensity of GDP by 2030 (relative to levels in 2005) 	<ul style="list-style-type: none"> Forestry emissions included in base year emissions intensity calculation Good Practice Guidance LULUCF of 2003 used to calculate emissions The inclusion of "non-forest land (cropland, grassland, wetlands and settlement)" will be determined at a later stage 	<ul style="list-style-type: none"> Malaysia points to "LULUCF legacy issues": past degradation including of peatlands Protection of coastlines is an adaptation priority: Shoreline Management Plans, National Coastal Vulnerability Index to sea-level rise in development
Philippines	<ul style="list-style-type: none"> "About 70%" of emission reductions by 2030 relative to the business-as-usual (BAU) scenario All sectors are included Target is conditional on international support 	<ul style="list-style-type: none"> 2006 IPCC Guidelines used as well as "Agriculture and Land Use (ALU) Software for agriculture, forestry and other land uses" 	<ul style="list-style-type: none"> Philippines engages in REDD+ "Marine ecosystems can play a crucial role with its potential on blue carbon" Declaration of 97 protected areas
RO Korea	<ul style="list-style-type: none"> Reduce emissions by 37% (or 850 MMt CO₂eq.) from BAU across sectors by 2030 (but see on LULUCF in next column) 	<ul style="list-style-type: none"> Inclusion of LULUCF to be decided later 	<ul style="list-style-type: none"> Korea "will partly use carbon credits from international market mechanisms to achieve its 2030 mitigation target"
Singapore	<ul style="list-style-type: none"> Reduce emissions intensity (GHG emissions per unit GDP) by 36% from 2005 levels in 2030 Achieve emissions peak (from which they will decrease) "around 2030" 	<ul style="list-style-type: none"> Emissions concerning land use change and forestry (LUCF) are accounted for 	<ul style="list-style-type: none"> "The array of natural ecosystems (including evergreen rainforest, mangroves, freshwater streams, freshwater swamp forest, coral reefs and mudflats) will continue to be conserved, with targeted programs for habitat enhancement."

Table 6. (I)NDC approaches of PEMSEA partner countries. (cont.)

Country	INDC/NDC Commitment	LULUCF Accounting	Policies, Measures & Mechanisms (LU-LUCF)
Thailand	<ul style="list-style-type: none"> • Reduce, unconditionally, GHG emissions by 20% by 2030 compared with business-as-usual (BAU) • With international support, this could be as much as 25% • Economy-wide, but see LULUCF 	<ul style="list-style-type: none"> • Inclusion of LULUCF to be decided later • Acknowledges vulnerability to coastline flooding 	<ul style="list-style-type: none"> • Climate change market mechanisms are encouraged • Thailand considers REDD+ engagement • To increase national forest cover to 40%, including “in particular, headwater and mangrove forests” • To develop participatory, integrated marine conservation and coastal rehabilitation plan to protect marine ecosystems and enhance climate proofing infrastructure to strengthen coastal protection against erosion
Timor-Leste	<ul style="list-style-type: none"> • No INDC/NDC submitted yet 	<ul style="list-style-type: none"> • The initial National Communication (NC) notes that the country holds “many coastal and marine resources including fish, seagrasses, seaweeds, coral reefs, mangrove forests and beaches...” • Initial NC includes explicit inclusion of mangroves as forest (primary or secondary) 	<ul style="list-style-type: none"> • Initial NC mentions 400 hectares of mangrove restoration on “swampy shrubland areas” as a “sink enhancement program” • The list of “key adaptation actions” includes the “protection and rehabilitation of mangrove ecosystems in priority areas to protect economic, social and environmental assets against climate risks”
Viet Nam	<ul style="list-style-type: none"> • Reduce, unconditionally, emissions across sectors by 8%, with the following breakdown <ul style="list-style-type: none"> ◦ Energy intensity per unit of GDP to be reduced by 20% compared to 2010 levels ◦ Forest cover will increase to the level of 45% • Reduction will go up to 25% on the condition of international support (30% energy intensity) 	<ul style="list-style-type: none"> • LULUCF covered, namely forest land, cropland, grassland, wetlands, settlements, other land • IPCC Guidelines are mentioned in general terms 	<ul style="list-style-type: none"> • REDD+ • NAMAs • Market experience (CDM) • Use of new market mechanisms are encouraged • Aims to “protect, restore, plant and improve the quality of coastal forests, including mangroves, especially in coastal estuaries and the Mekong and Red River deltas” • Plans to increase the area of protection forest in coastal areas to 380,000 hectares, including 20,000 – 50,000 ha of additional mangrove planting • To implement integrated coastal zone management • To use sea level rise scenarios in urban and land use planning for infrastructure, industrial parks, coastal and island resettlement areas • To implement anti-inundation measures for large coastal cities • To protect the coastline and riverbanks

Moving Forward Under NDCs in the Region

The Paris Agreement foresees various mechanisms for the implementation and gradual strengthening of country NDCs. At the GHG inventory level, procedures have been set in place to improve, over time, both the completeness and the accurateness of the data.⁹⁵ Given the recent publication of the IPCC Wetland Supplement,⁹⁶ it should be possible to cover coastal wetland related removals and emissions within the territorial waters of countries.⁹⁷ The institutionalization of revision cycles for biennial reports (by industrialized countries) and biennial update reports (by developing countries) will further strengthen the reporting standards (if not in itself the quality of the data).

Other than improving completeness and accuracy, countries are under an obligation to strengthen their mitigation commitments over time. The Paris Agreement includes a dynamic mechanism to that effect (also referred to as the “Ratchet Mechanism”) that triggers procedural milestones to assess and improve a country’s mitigation ambition over time.⁹⁸ Each Party must prepare, communicate and maintain successive NDCs (Art. 14.2 Paris Agreement). Every five years, countries must update their NDC (Art. 4.9). Each time, the update needs to represent “a progres-

sion beyond the Party’s then current nationally determined contribution” (Art. 4.3), and it needs to consider the five-yearly “*global stocktake*” exercise (the first to happen in 2023), mandated under Article 14 as an assessment of “*the collective progress towards achieving the purpose of this Agreement and its long-term goals*” (ibid.) Note that while countries determine their NDC individually, without assuming a legal obligation or liability vis-à-vis the result, they do have an obligation to pursue mitigation actions with the “*aim of achieving the objectives*” of their NDCs (Art. 4.2).

Countries do not have to do it alone. Rather, the Paris Agreement sets incentives for bilateral and multilateral cooperation, including through emissions trading. The point of departure is what is called “voluntary cooperation” of countries to allow for higher mitigation and adaptation ambition and the promotion of “*sustainable development and environmental integrity*” (Art. 6.1). Article 6 includes three cooperation formats:⁹⁹

- **Cooperative Approaches.** Parties may engage in “*voluntary cooperation*” (Art. 6.1) and “*cooperative approaches*”, using “*internationally transferred mitigation outcomes*” (Art. 6.2), a terminology that is broad enough to allow for different types of trades. It is noted that trades can cover all sectors, including sequestration (removals by sink).¹⁰⁰

⁹⁵ The Paris Agreement and the decision accompanying it (the Paris Decision) lay out several provisions, including: Parties must provide their NDCs with ‘*necessary*’ levels of ‘*clarity, transparency and understanding*’ (Paris Decision, para. 13) in accordance with what has been decided under the Paris Decision and what will be decided by the CPA. Furthermore, all Parties shall account for their actions and results achieved in ‘*implementing the NDCs*’ (Art. 4.13 Paris Agreement). In accounting for emissions and removals ‘*corresponding to*’ their NDCs, Parties shall ‘*promote environmental integrity, transparency, accuracy, completeness, comparability and consistency, and ensure the avoidance of double-counting*’, in accordance with guidance to be adopted (ibid).

⁹⁶ Hiraishi, Takahiko, et al. 2014. 2013 supplement to the 2006 IPCC guidelines for national greenhouse gas inventories: Wetlands.

⁹⁷ *Ibid.* (op. cit.), Cross-Cutting Issues and Reporting: “Coastal wetlands can occur in any of the six IPCC land-use categories but also in coastal areas that are not part of the total land area of the country. For example, a mangrove wetland with trees may be classified as Forest Land, a tidal marsh used for grazing may be classified as Grassland, and a seagrass meadow used for aquaculture may be classified as Settlements. Emissions/removals from coastal wetlands that are not part of the total land area (e.g. seagrass meadows) should be reported separately and the associated areas excluded from the total land area and from the land-use example, forest management activities in mangroves classified as Forest Land may need to be split between areas included in the total land area and not included in the total land area. In reporting the emissions/removals from mangrove forest management activities, emissions/removals from both areas would be reported under Forest Land, but only the land areas of the mangroves included in the total land area would be included in the total Forest Land areas and reported in the land area matrix. The classifications of coastal wetlands are country-specific, but in all cases appropriate subcategories should be used in the reporting to reflect the specific land use and management as well as to indicate whether the emissions come from areas included or excluded from the total land area of the country...”

⁹⁸ For details see C. Streck et al. 2016. The Paris Agreement: A Beginning. *Journal for European Environmental & Planning Law*, 133-29.

⁹⁹ For details see Streck et al. (op. cit.).

¹⁰⁰ Paris Decision, para 37.

- **Sustainable Development Mechanism (*name still to be formally adopted*)**. Based on a last-minute intervention by several countries, the Paris Agreement defines a mechanism that allows private and public entities to support mitigation projects that generate transferrable GHG emissions (Art. 6.4). Programs and projects—the Paris Agreement avoids using either term—developed under this new mechanism can generate “*emission reductions*” that may be used by another Party to “*fulfill*” its NDC. The mechanism is implemented under the “*authority and guidance*” of the main institutional body of the Paris Agreement, which, according to the Paris Decision, is intended to develop relevant “*modalities and procedures*.”¹⁰¹ The provision in the Paris Decision links back to the mechanisms of the Kyoto Protocol, namely the CDM, when requesting that the new mechanism be built on their experience (para 38.f). Similar to the CDM, the mechanism addresses subnational public and private entities, and it foresees a “*share of proceeds*” to cover both administrative costs and adaptation needs for nations most vulnerable to climate change (Article 6.6). This opens a future for the Adaptation Fund, created under the Kyoto Protocol. However, unlike the CDM, the new mechanism appears open to all sectors, including blue carbon. The main threshold is that it must “*deliver an overall mitigation in global emissions*” (Art. 6.4.d), that is, it must go beyond offsetting and have a net positive mitigation effect. Also, emission reductions may be accounted for only once in the context of
- NDCs, either by the host Party or by another Party (Article 6.5).
- **Framework for non-market approaches**. The Paris Agreement recognizes “*the importance of integrated, holistic and balanced non-market approaches*” (Art. 6.8) to assist Parties with implementing their NDCs, in the context of sustainable development and poverty eradication. It aims at both mitigation and adaptation, “*enhance[s] public and private sector participation*” and seeks opportunities for coordination “*across instruments and relevant institutional arrangements*.” The conceptual scope and meaning of non-market approaches, as opposed to instruments seen as (if no longer called) market mechanisms and for which we find precedence in the Kyoto mechanisms, is hard to gauge. In a technical paper from 2014, the UNFCCC secretariat summarized non-market approaches as “*any actions that drive cost-effective mitigation without relying on market-based approaches or mechanisms* (i.e. without resulting in transferable or tradable units).”¹⁰² The technical paper listed as examples, from country experience, fiscal instruments (such as carbon taxes) and regulation, but also voluntary agreements on mitigation action and results-based payments for REDD+. The concept, in this interpretation, is very wide, and there will be much work ahead for SBSTA,¹⁰³ which is charged with preparing a draft work programme until next year’s session.¹⁰⁴

¹⁰¹ Paris Decision, para 37 and 38 (the latter provision involving SBSTA).

¹⁰² UNFCCC Secretariat. “Non-market based approaches: Technical Paper, FCCC/TP/2014/10, 24 November 2014.” <http://unfccc.int/resource/docs/2014/tp/10.pdf>. Accessed 2017 February 22.

¹⁰³ UNFCCC. [n.d.]. “Subsidiary Body for Scientific and Technological Advice (SBSTA).” <http://unfccc.int/bodies/body/6399.php>. Accessed 2017 February 23.

¹⁰⁴ Paris Decision, paras 40 and 41.

Policy Opportunities for PEMSEA Partner Countries

While all PEMSEA partner countries in their emerging climate change architecture put some emphasis on LULUCF and the importance of forests generally, there are substantial differences in the approach to blue carbon ecosystems and the readiness to engage in both adaptation and mitigation mechanisms.

Cambodia, China, Japan and Viet Nam formulate forest-sector related targets, while five other countries (DPR Korea, Indonesia, Malaysia, the Philippines and Singapore) include LULUCF emissions or LUCF emissions (excluding emissions from unchanged land use) as part of their cross-sectoral targets. RO Korea and Thailand will decide on the inclusion of LULUCF in their NDCs at a later stage.

The importance of conserving and/or restoring natural ecosystems—often specifically referring to coastal ecosystems and mangroves—is highlighted in all (I)NDCs from the region. While the policy perspective is mostly one of adaptation, several PEMSEA countries make specific reference to forest *mitigation* activities, in the form of REDD+ (Cambodia, Indonesia, the Philippines, Thailand) or other interventions, namely afforestation, reforestation or restoration (Viet Nam and Timor-Leste).¹⁰⁵

Several PEMSEA countries have been strong performers in early emissions trading (especially China and Viet Nam, but also Thailand, the Philippines and Japan), and several of them have established emissions trading schemes of their own (China, RO Korea and Japan) or contemplate some form of trading (Thailand). The Philippines explicitly mentions the “blue carbon” potential, and Viet Nam states its strong

interest in developing future market mechanisms, while highlighting the need to protect, restore and improve coastal forests. Both Japan and RO Korea indicate that they may invest in forest carbon emissions projects or programs to offset parts of their national emissions.

Overall, the analysis points to several blue carbon policy and finance opportunities for PEMSEA partner countries:

1. As climate finance support for recipient countries is increasingly *guided and shaped* by the countries’ NDCs, the fact that coastal ecosystems are featured prominently makes blue carbon interventions a clear finance priority.
2. A range of international climate finance programs—from the Green Climate Fund to programs of international finance institutions to the multi-donor NAMA Facility to single donor platforms—require *official vetting* by the recipient country government before an intervention is approved.¹⁰⁶ An NDC reference facilitates official vetting, just as the failure to be mentioned in a country’s NDC lowers the odds for an intervention type to be approved.
3. Some NDCs are drafted in a way that highlights linkages and interconnectivities between mitigation efforts, adaptation efforts and cross-sectoral approaches, such as coastal planning. Viet Nam, for instance, spelled out what can be seen as a detailed roadmap for cross-sectoral engagement. The text commits to “protect, restore, plant and improve the quality of coastal forests, including mangroves, especially in coastal estuaries and the Mekong and Red River deltas”. Viet Nam plans to increase the protection of forest in coastal areas to 380,000 ha, including 20,000–50,000 ha of additional mangrove

¹⁰⁵ Information gathered from the initial NC.

¹⁰⁶ For an overview of climate finance programs and funding platforms see www.ndcpartnership.org.

planting; implement integrated coastal zone management; and it proclaims to work with sea level rise scenarios in urban and land use planning for infrastructure, industrial parks and coastal and island resettlement areas, to implement anti-inundation measures for large coastal cities and generally to protect the coastline and riverbanks. Such cross-cutting methodology makes NDCs powerful documents to inspire projects and programs and to secure their comprehensive design.¹⁰⁷

4. Blue carbon interventions, whether included in all-country REDD+ implementation or carried forward in the form of projects, are driven by local communities, and above all, they involve the improvement of natural environments and livelihoods alike. This makes them strong contenders for any climate funds offered and creates additional value on voluntary carbon markets.
5. Several PEMSEA countries are particularly open to carbon crediting mechanisms—the Philippines and Viet Nam on the side of beneficiaries, and Japan and potentially RO Korea on the side of investor countries. While new mechanisms are still in the design phase, countries have been given a green light to engage bilaterally in transactions, and the group of PEMSEA countries, with PEMSEA, CI and TNC perhaps providing the institutional framework (see next point), are well positioned to take the lead. Japan already has experience with bilateral REDD+ projects, so the design of a dedicated blue carbon project or program would not need to start from scratch.
6. PEMSEA, CI, TNC, the Blue Carbon Initiative and other similar organizations are in the unique position to

advise PEMSEA countries on potential intervention formats, to collect and streamline country initiatives and cross-country experience and to operate as focal point and functional interface for regional initiatives. The Partnership for Blue Carbon, supported by the Australian government, is another potentially important partner in such an activity. This role could extend to the facilitation of bilateral and multilateral initiatives under Article 6.2 of the Paris Agreement: between potential beneficiary and investor countries, but also with a view of developing a “Regional Blue Carbon Facility”, which could ultimately act as a trading platform and clearinghouse among PEMSEA country partners and other countries seeking to engage in blue carbon initiatives in the region.

Difficulties remain: as highlighted in this report, carbon stock, emissions and removals data for blue carbon ecosystems remains incomplete. Guidelines for inventories, monitoring, accounting, etc. are not readily available. And relevant blue carbon ecosystems, notably seagrasses, remain outside any reporting, accounting or NDC framework.

However, this is likely to change, and the new set of NDCs is likely to address blue carbon ecosystems with more precision and comprehensive rigor. Regional and global organizations and initiatives, such as PEMSEA, the Blue Carbon Initiative, the Partnership for Blue Carbon and others, can help to facilitate further change.

Table 7 highlights potential priority areas of engagement for PEMSEA partner countries.

¹⁰⁷ For an example see UNDP’s project for Viet Nam (“Improving the resilience of vulnerable coastal communities to climate change related impacts in Viet Nam”), which includes mitigation and adaptation and urban planning activities.

Table 7. Potential priority areas of engagement for PEMSEA partner countries.

Country	Priority Areas/Recommended Next Steps
Brunei Darussalam	<ul style="list-style-type: none"> • Prepare a complete blue carbon inventory (including seagrasses) • Identify instances of degradation (if any), potential drivers and mitigation strategies
Cambodia	<ul style="list-style-type: none"> • Acknowledge mangroves as a mitigation opportunity • Build a mangrove focus into the REDD+ strategy • Identify an area for a mangrove-based NAMA
China	<ul style="list-style-type: none"> • Build mangrove and other blue carbon ecosystems into a nation-wide inventory • Create a blue carbon project facility to generate projects, modelled on the CDM and VCS, as offsets for the domestic carbon markets
DPR Korea	<ul style="list-style-type: none"> • Identify blue carbon habitats across the country • Start building inventory-level datasets • Develop a cross-sectoral NAMA involving blue carbon ecosystem conservation and restoration, cookstoves, off-grid solar and fisheries
Indonesia	<ul style="list-style-type: none"> • Build capacity on quantification of blue carbon stocks and stock change • Include blue carbon in national reporting of GHG emissions and sinks • Support interagency coordination on coastal ecosystem conservation • Integrate a blue carbon target both in the conditional and the unconditional target • Encourage pilots of community-based blue carbon projects by giving out no-cost coastal and mangrove habitat management licenses
Japan	<ul style="list-style-type: none"> • Focus on blue carbon in bilateral and regional engagements • Include blue carbon projects in the JCM mechanism • Undertake feasibility assessment of blue carbon based cooperation within the scope of Article 6.2 Paris Agreement
Malaysia	<ul style="list-style-type: none"> • Build capacity on quantification of blue carbon stocks and stock change • Support interagency coordination on coastal ecosystem conservation • Include blue carbon in national reporting of GHG emissions and sinks • Prepare the adoption of an absolute mitigation target for the blue carbon sector
Philippines	<ul style="list-style-type: none"> • Build capacity on quantification of blue carbon stocks and stock change • Support interagency coordination on coastal ecosystem conservation • Include blue carbon in national reporting of GHG emissions and sinks • Work through national Blue Carbon Working Group to develop a national blue carbon workplan • Pilot blue carbon projects and seek international cooperation on trading blue-carbon related emissions
RO Korea	<ul style="list-style-type: none"> • Identify blue carbon ecosystems around the country • Include blue carbon in national reporting of GHG emissions and sinks • Link domestic and regional blue carbon projects as offsets to the domestic emissions trading scheme
Singapore	<ul style="list-style-type: none"> • Facilitate cross-boundary blue carbon interventions • Set up a blue carbon trading infrastructure for the region
Thailand	<ul style="list-style-type: none"> • Build capacity on quantification of blue carbon stocks and stock change • Include blue carbon in national reporting of GHG emissions and sinks • Support interagency coordination on coastal ecosystem conservation • Link blue carbon action to the Partnership for Market Readiness and to emerging emissions trading tools • Build a "Blue Carbon NAMA" targeting the aquaculture industry
Timor Leste	<ul style="list-style-type: none"> • Build capacity on quantification of blue carbon stocks and stock change • Include blue carbon in national reporting of GHG emissions and sinks • Support interagency coordination on coastal ecosystem conservation • Build the NDC to integrate and focus on blue carbon environments
Viet Nam	<ul style="list-style-type: none"> • Build capacity on quantification of blue carbon stocks and stock change • Include blue carbon in national reporting of GHG emissions and sinks • Support interagency coordination on coastal ecosystem conservation • Include a "Blue-Carbon-And..." window in NAMA infrastructure ("Blue Carbon and Shrimp Industry", "Blue Carbon and Rice Paddies", "Blue Carbon and Fisheries", etc.) • Link blue carbon action and Partnership for Market Readiness • Actively seek cooperation on Article 6.2 Paris Agreement with interventions focusing on domestic blue carbon habitats

Linking Climate Finance and Blue Carbon Investment

The various NAMA initiatives point to an emerging pattern of blue-carbon related investments linking conservation and restoration aims—climate change mitigation and adaptation goals—with activities that promise a return-on-investment. These activities may be related to coastal-based *commodities* such as fisheries, aquaculture and ocean-based energy, or *services* such as habitat resilience and flood prevention (thus, flood insurance schemes), tourism or emissions trading and biodiversity compensation schemes. Or they involve measures with an indirect effect on habitats (e.g., energy-efficient cookstoves that lower the pressure on coastal timber or off-grid energy production, which adds economic opportunities and may reduce pressure on land).

Other than seeking dual benefits (i.e., conservation and economic opportunities), this coupling presents an important opportunity to attract private capital in an environment that historically was funded almost exclusively by public donors (aid and development) or philanthropists. Natural capital investments are on the rise, with the international climate change architecture acting as an important catalyst. Blue carbon inventories and monitoring systems have become highly regarded proxies for ecosystem evaluation, facilitated by deployment of state-of-the-art technology to track land use change across landscapes (e.g. satellite-based). REDD+ strategy development land tenure assessments provide blueprints for due diligence assessments for land development at large. Public climate finance sources offer senior debt and mezzanine instruments¹⁰⁸ to allow on-lending by local financial institutions in the agriculture or fisheries sector, and climate micro-finance supported by public climate finance has become an industry in its own right.

Focused blue carbon engagement allows the PEMSEA countries to bundle scientific, technical, legal and financial actions using existing knowledge facilities, financial support systems and practical blueprints. To be sure, the climate architecture under the Paris Agreement does not offer an automated system of capacity-development, strategic advice and finance. Rather, true to the methodological paradigm of bottom-up action, it falls to countries and their individual ambition to build their climate change intervention portfolio.

For a pro-active government, the blue carbon environment holds considerable momentum. **Table 8** lists several public funds that could directly finance conservation and restoration activities in mangroves and other blue carbon ecosystems. Beyond these public funds, however, there is a range of knowledge and funding tools that could be used to enhance and invest in a country's natural capital. The Climate Risk Early Warning System (CREWS), for instance, embedded in the Global Facility for Disaster Reduction and Recovery (GFDRR), aims at approving meteorological warning systems. A country may apply to this program and link the supported measure to the development of a mangrove-based weather protection response system. If it also traces human-induced carbon fluctuations, it can add a carbon crediting component.

Similar cross-sectoral combinations are possible at the project level. A marine protected area, for instance, that promotes healthy mangrove forests as a means to improve fisheries, should consider developing a carbon project to achieve co-funding. At a higher national or regional level, again, these efforts can be helped by offering crediting mechanisms (REDD+ implementation, say, or a dedicated blue carbon regional mechanism) to project developers, and by setting up institutional structures (such as Roundtables) that help influence drivers of deforestation beyond the reach of the project level (e.g., supply chains).

¹⁰⁸ A hybrid of debt and equity financing that gives the lender the rights to convert to an ownership or equity interest in the company in case of default, after venture capital companies and other senior lenders are paid.

Table 8. Public Climate Funds and Facilities Available for PEMSEA Countries (REDD+ and Blue Carbon Focus) – Examples (bilateral and multilateral)¹⁰⁹

Fund	Governance	Special Focus
Adaptation Fund	Adaptation Fund Board (administered by World Bank)	Adaptation
BioCarbon Fund	World Bank	Sustainable Forest Landscapes
Forest Carbon Partnership Facility	World Bank	Participating PEMSEA countries: <ul style="list-style-type: none"> • Cambodia • Indonesia • Thailand • Viet Nam
Forest Investment Program (Strategic Climate Fund)	World Bank (with, among others, Asian Development Bank)	REDD+ support
Global Environment Facility Trust Fund	Global Environment Facility	Adaptation, Mitigation
Global Climate Change Alliance	European Commission	<ul style="list-style-type: none"> • Focus on Least Developed Countries (LDCs) and Small Island States (SIDS) • Seeks to enhance regional partnerships, in Asia currently the Mekong River Commission
Green Climate Fund	Green Climate Fund Board	<ul style="list-style-type: none"> • Mitigation, Adaptation, REDD+
Global Facility for Disaster Reduction and Recovery	World Bank	<ul style="list-style-type: none"> • Supports risk management projects
Global Resilience Partnership	Rockefeller Foundation, USAID, Sida	<ul style="list-style-type: none"> • One of three focus regions is South and South East Asia
International Climate Initiative	Germany	<ul style="list-style-type: none"> • Mitigation, Adaptation, REDD
International Forest Carbon Initiative	Australia	<ul style="list-style-type: none"> • Currently supports Indonesia
International Forest Climate Initiative	Norway	<ul style="list-style-type: none"> • Indonesia is currently funded under the initiative
Joint Crediting Mechanism	Japan	Currently operates in 16 countries, among them: <ul style="list-style-type: none"> • Cambodia • Indonesia • Thailand • Philippines • Viet Nam
Least Developed Country Fund	World Bank	<ul style="list-style-type: none"> • Cambodia • Timor-Leste
NAMA Facility	Germany, UK	<ul style="list-style-type: none"> • Innovative climate finance approaches
Pilot Program for Climate Resilience (Strategic Climate Fund)	World Bank	Currently funded: <ul style="list-style-type: none"> • Cambodia • Philippines
Special Climate Change Fund	GEF	<ul style="list-style-type: none"> • Adaptation and Technology Transfer
UN REDD	UNDP-FAO	<ul style="list-style-type: none"> • REDD+ readiness

¹⁰⁹ Excludes Japan and for most purposes the Republic of Korea.



Photo by SCA/Crooks

5 Practical Steps to Advance Blue Carbon Interventions

The countries of East Asia, indeed coastal countries around the world, can enact or strengthen actions that improve management of blue carbon ecosystems either to support climate change policy responses and interventions, as described in this report, or for other purposes. Conservation and restoration of blue carbon ecosystems (and other coastal and ocean ecosystems) cuts across and underpins many aspects of “blue economy” sectors. Improved management also benefits from regional and cross-border collaboration. **Table 9** below highlights a framework of actions that

PEMSEA countries can take to advance the management of blue carbon ecosystems, climate response planning and blue economy growth.¹¹⁰ The framework is based upon three main pillars: 1) awareness building, 2) knowledge exchange and 3) acceleration of practical action, including making use of emerging climate change instruments. While some countries may already be advanced in some of these actions, there is value in regional comparability. PEMSEA, CI and TNC have an opportunity to help facilitate communication and capacity building as partner countries enact relevant aspects of this framework.

Table 9. Recommendations for countries to incorporate blue carbon ecosystems into integrated coastal management, climate response, biodiversity conservation and blue economy planning.

Action	Benefit	Actor
Building Awareness		
Include blue carbon in policy dialogue.	Supports development of national and subnational policies, cooperation between governments and intra-government agencies and inclusion of private sector and community groups.	National government; International agencies; International NGOs; Academic community.
Apply 2013 IPCC Wetland Supplement and include blue carbon ecosystems in GHG National Inventory and Communications.	Improved quantification of emissions and removals due to land management. Enables setting of goals and benchmarks for management plans.	National Government.
Report trends of coastal ecosystems, including improved mapping of blue carbon ecosystems, their change through time, threats and status.	Supports management planning and inclusion of blue carbon ecosystems in GHG national Inventories and communications.	National government; International agencies; Academic community.

¹¹⁰ See the Blue Carbon Partnership for the international application of this framework structure <http://bluecarbonpartnership.org/>

Table 9. Recommendations for countries to incorporate blue carbon ecosystems into integrated coastal management, climate response, biodiversity conservation and blue economy planning. (cont.)

Action	Benefit	Actor
Facilitate Knowledge Exchange		
Join networks such as the International Partnership for Blue Carbon¹¹¹ and the International Blue Carbon Initiative.¹¹²	Bring together key organizations to coordinate international activities.	National government; International NGOs, Academic Community.
Facilitate / contribute to technical and policy workshops (e.g., The Blue Carbon Initiative).¹¹³	Enable communication between technical experts and shared science, policy and implementation experience.	National government; International NGOs; Private sector; Academic community.
Support science programs and technical analysis.	Improved quantification of blue carbon benefits and understanding of intervention opportunities.	National government; International NGOs; Private sector; Academic community.
Develop knowledge products and demonstration activities, e.g., activities under GEF Blue Forest Project¹¹⁴ and by Restore America's Estuaries.¹¹⁵	Demonstration and communication of experience and good practice to support mainstreaming and upscaling of blue carbon interventions.	National government; International NGOs; Private sector; Academic community.
Accelerate Practical Action		
Investigate appropriate policy frameworks for including blue carbon ecosystems within national commitments to the Paris Agreement.	Including blue carbon ecosystems within NDCs and related plans provides guidance to coastal planners and assists in securing international funding for climate adaptation and mitigation.	National government.
Include management of blue carbon ecosystems within integrated coastal management plans.	Integrated coastal management plans help to steer on-the-ground climate response and blue economy development. Including the status of, and goals of for, blue carbon ecosystems can provide a foundation for broader coastal management.	National and local government.
Assess and promote national opportunities for conservation and restoration of blue carbon ecosystems, including quantification of GHG benefits.	Blue carbon ecosystems are being lost across East Asia at a high rate. Reversing these losses support components of NDCs, the UN SDGs and blue economy growth.	National and local government.
Provide training and technical support to local and national government agencies, field schools and communities on the value of blue carbon ecosystems and good practice for conservation and restoration.	Experience in restoring blue carbon ecosystems exists, but success rates are still relatively low. Training and improved planning can support more successful delivery.	International development organizations; National government; International NGOs; Private sector.

¹¹¹ International Partnership for Blue Carbon. 2017. <http://bluecarbonpartnership.org/>.

¹¹² *Ibid.*

¹¹³ *Ibid.*

¹¹⁴ GEF Blue Forests. 2016. <http://www.gefblueforests.org/>.

¹¹⁵ Restore America's Estuaries. 2016. "Blue Carbon Resources." <https://www.estuaries.org/bluecarbon-resources>.

Table 9. Recommendations for countries to incorporate blue carbon ecosystems into integrated coastal management, climate response, biodiversity conservation and blue economy planning. (cont.)

Action	Benefit	Actor
Develop climate change adaptation strategies that consider migration of blue carbon ecosystems with sea level rise and human impacts (such as dam construction) on sediment supply to coastal regions.	Space is one of the scarcest resources in coastal areas. Adapting to climate change requires that plans incorporate landward movement of coastal assets including blue carbon ecosystems. There is an opportunity to plan buffer areas of no or low development that will both create space for coastal wetlands to migrate landwards in the future as well as reduce risk to coastal communities from climate change	National and local government.
Include blue carbon ecosystems in coastal vulnerability assessments.	Along with hard infrastructure, natural infrastructure, including blue carbon ecosystems, is an important element in reducing ecosystem and human vulnerability to climate change. Developing blue carbon vulnerability assessments will empower governments and communities to manage natural resources into the future.	International development organizations; National government; International NGOs; Private sector.
Include blue carbon ecosystems in national economic development plans.	Recognizing the natural capital value of intact and restored blue carbon ecosystems in economic development plans can support development of sustainable blue economies.	National and local government.
Include blue carbon ecosystems as a component of natural infrastructure.	Coastal and river wetlands provide valuable flood risk reduction services. Including wetlands in development plans provides additional levels of protection during storm and high-low events, along with additional ecosystem services not provided by hard infrastructure.	International development organizations; National government; International NGOs; Private sector.
Include blue carbon ecosystems within marine protected areas.	Blue carbon ecosystems are important elements of marine protected areas, supporting biodiversity, providing fish nurseries and other services underpinning marine ecology and productivity. Agreements established to support MPAs provide a basis for other blue carbon interventions.	International development organizations; National government; International NGOs.
Include blue carbon ecosystems as part of marine spatial planning and other tools for managing multi-use coastal landscapes.	Marine spatial planning offers the opportunity to map and track changes in blue carbon ecosystems through time and to support alignment of management approaches for their conservation.	International development organizations; National government; International NGOs; Private sector.

Table 9. Recommendations for countries to incorporate blue carbon ecosystems into integrated coastal management, climate response, biodiversity conservation and blue economy planning. (cont.)

Action	Benefit	Actor
Develop/apply soil management plans for watershed and coastal regions.	Improved soil management results in reduced release of carbon either through erosion or directly to the atmosphere in the form of carbon dioxide or methane.	National and local government.
Correlate health of blue carbon ecosystems with industry inputs and outputs of blue economy.	Clarify the interdependency of blue economy industries with function of coastal ecosystems. Minimize industry environmental liabilities and maximize benefits.	National and Local government; Private sector.



Photo by SCA/Crooks

6

Addendum: Blue Carbon Ecosystems as Part of an East Asia Blue Economy

While the economic perspectives of blue carbon ecosystems were beyond the scope of this report, the following addendum provides brief background and reference concepts to facilitate the inclusion of “blue economy” in the follow-on dialogue and activities around blue carbon by PEMSEA and its partners.

Private Investment for Conserving and Restoring Blue Carbon Ecosystems

PEMSEA’s 2015 report on *Blue Economy for Business in East Asia* revealed that coastal restoration, protection and climate change adaptation are among the sustainable development investment priorities for companies surveyed in the region, over a 3 to 5-year time horizon. Blue carbon ecosystems must undoubtedly be factored in as prominent elements of these investments.

Effective approaches for securing private investment and related expertise is critical, as a significant gap exists globally and regionally between the need for conservation and climate related investments and the availability of public and philanthropic funding. The Coalition for Private Investment in Conservation (CPIC)—an effort launched by

Credit Suisse, The Nature Conservancy, International Union for Conservation of Nature (IUCN) and Cornell University¹¹⁶ to develop new investment models to help close the current conservation funding gap—provides the following perspective on the situation:

*In order to deliver the volume of investment needed to address the scale of conservation challenges, investment opportunities that provide measurable, science-based conservation benefits and social impact to participating communities and to biodiversity, while delivering at-scale financial returns for investors, will be necessary.*¹¹⁷

Furthermore, finding innovative ways to secure private investment is a particularly important challenge for protection and restoration of blue carbon ecosystems, as noted in the Coastal Blue Carbon report from IUCN:

*Finding appropriate funding sources to set up a coastal wetland carbon project or develop a national carbon program (which includes or is solely focused on coastal wetlands) is often a challenge. Additionally, carbon finance alone often cannot support the necessary management activities.*¹¹⁸

¹¹⁶ Others members include (but are not limited to) the Global Environment Facility, Encourage Capital, WWF, the European Investment Bank, Convergence, Conservation Finance Alliance, Conservation Finance Network, Global Ocean Trust and World Forum on Natural Capital.

¹¹⁷ Coalition for Private Investment in Conservation. “Statement of Intent accessible.” <http://www.conservationfinancenetwork.org/2016/09/28/coalition-launched-to-scale-up-conservation-finance>.

¹¹⁸ Herr, D. T. et al. 2015. Coastal “blue” carbon. A revised guide to supporting coastal wetland programs and projects using climate finance and other financial mechanisms. IUCN, Gland. https://www.iucn.org/sites/dev/files/import/downloads/wetlands_carbon_finance_paper_final_lr.pdf. Accessed 2017 January 25.

The emerging fields of conservation investment and blended finance, both independently and working together, offer approaches for securing private investment, and in some cases may be able to augment public and philanthropic sources of carbon and/or climate funding.

Conservation Investment

Conservation Investments are investments intended to return principal or generate profit while also resulting in a positive impact on natural resources and ecosystems. In addition, conservation impacts must be the intended motivation for making the investment; they cannot be simply a by-product of an investment made solely for financial return.¹¹⁹

Conservation Investment is considered a segment of the wider Impact Investment market and is growing substantially. TNC finds in its 2016 *Investing in Conservation* report that in just two years (2013-2015), the total private capital committed to conservation investments jumped by 62%, to a total committed private capital of \$8.2 billion USD tracked from 2004 to 2015.¹²⁰

In its report *Conservation Finance: Moving Beyond Donor Funding Toward an Investor-driven Approach*, Credit Suisse finds that “investors say they are looking for investable projects with clear investment characteristics, run by managers or trusted funds with conservation and financing experience and track record.”¹²¹ These can include:

1. Investments in underlying ecosystems with the goal of capital protection. *The acquisition of forests, freshwater or deserts, or usage rights tied to a long-term conservation commitment only make sense from a financial perspective if they generate a financial return and thereby become actual financial assets (though the underlying ecosystem may have intrinsic value for some investors or generate other benefits, such as tax breaks).*

2. Investments in infrastructure and sustainable management of ecosystem services to achieve financial returns.

These investments can create economic value under the constraint of conservation, for example with lodges and trails to foster ecotourism or solar arrays for power generation. Ecosystem services (such as watershed protection) and goods derived from sustainable forestry, agriculture, or aquaculture can also provide cash flows. Such cash flows often depend on regulatory requirements or industry certifications that support premium pricing.

3. Investments in ecosystem market and regulatory mechanisms to enhance returns.

These are financial instruments such as securities and derivatives, as well as corporate intermediaries leveraging regulatory requirements. Examples include voluntary or mandatory offsets, subsidized power production or permit and rights issuance and trading.

Blended Finance

The Organization for Economic Cooperation and Development (OECD) and World Economic Forum offer the following definition of Blended Finance:

The strategic use of development finance and philanthropic funds to mobilize private capital flows to emerging and frontier markets, characterized by:

- *Leverage: Use of development finance and philanthropic funds to attract private capital.*
- *Impact: Investments that drive social, environmental and economic progress.*
- *Returns: Returns for private investors in line with market expectations based on perceived risk.*

*Blended Finance can be used across a range of structures, geographies and sectors using a range of instruments. Deals bring together different stakeholders that partner in a fund or transaction, involving a mixture of development funding and private investors.*¹²²

¹¹⁹ J.P. Morgan. “State of Private Investment in Conservation 2016 A Landscape Assessment of an Emerging Market.” <https://www.cbd.int/financial/privatesector/g-private-wwf.pdf>. Accessed 2017 February 22.

¹²⁰ *Ibid.*

¹²¹ Credit Suisse. 2014. “Conservation Finance: Moving Beyond Donor Funding Toward an Investor-driven Approach.” <https://www.cbd.int/financial/privatesector/g-private-wwf.pdf>. Accessed 2017 February 22.

¹²² World Economic Forum and Organization for Economic Cooperation and Development. 2015. “A how-to guide for Blended Finance: a practical guide for Development Finance and Philanthropic Funders to integrate Blended Finance best practices into their organizations.” http://www3.weforum.org/docs/WEF_Blended_Finance_How_To_Guide.pdf. Accessed 2017 February 25.

As an example, The Nature Conservancy received a grant from Convergence Design Funding to develop a blended finance proof of concept grant in 2016. TNC will use the funding to design a Blue Bond that will finance debt conversions for Small Island Developing States that face climate change challenges. Debt conversions provide funding to governments in exchange for their commitment to improve policies and invest in marine conservation and climate adaptation activities.

Developing a substantial number of these types of investment deals in the region will likely require significant capacity building among a range of stakeholders over time. An opportunity exists for partners in the region to collaborate as a consortium in providing the regional leadership and expertise for developing innovative and systematic investment approaches as described above. This will be essential for attracting and securing the prioritized private investment in blue carbon ecosystems that can support the envisioned growth of a blue economy in East Asia.

Blue Carbon and the Blue Economy

From an integrated coastal management perspective, East Asia's blue carbon ecosystems are valued for their role as vital resources supporting the development of a blue economy in the region. As defined by PEMSEA partner countries in the 2012 Changwon Declaration, blue economy is "a practical ocean-based economic model using green infrastructure and technologies, innovative financing mechanisms and proactive institutional arrangements for meeting the twin goals of protecting our oceans and coasts and enhancing its potential contribution to sustainable development, including improving human well-being, and reducing environmental risks and ecological scarcities."¹²³

In this context, PEMSEA's *Blue Economy for Business in East Asia* report identifies four key elements of marine

and coastal economic activities that are considered blue economy:

1. Protects, restores and sustains healthy coastal and marine ecosystem services
2. Generates sustainable, equitable economic benefit and inclusive growth
3. Integrates approaches between multiple industries and government
4. Innovates, informed by the best science

Blue carbon ecosystems' role aligns most directly with the first of these elements. The rising strategic importance of ecosystems services and natural capital management in blue economy development planning is reflected in the following excerpt from the World Bank assessment *Toward a Blue Economy: a Promise for Sustainable Growth in the Caribbean*:

*"These drivers of change in the status of the Caribbean Sea's natural capital assets constitute an important constraint and significant risk to the potential growth of the region's ocean economy, similar to the risks to the global ocean economy. In some cases, the decline of these assets may prevent sectors and industries from reaching their potential in the region; for others, it may create regulatory uncertainty that also presents a significant risk."*¹²⁴

Research is underway by PEMSEA Country Partners to further develop a framework for understanding the blue economy in East Asia. The research will culminate in a set of National State of Oceans and Coasts Reports and a comprehensive *Regional Seas of East Asia State of Oceans and Coasts* report, planned for release in 2018 that will include:

- Definitions and methodologies for assessing blue economy
- The contribution of blue economy activities to national economies in East Asia, including the value of ecosystem services in coastal and ocean areas

¹²³ PEMSEA. 2015. "Blue Economy for business in East Asia: towards an integrated understanding of Blue Economy." <http://www.pemsea.org/publications/reports/blue-economy-business-east-asia-towards-integrated-understanding-blue-economy>. Accessed 2017 February 28.

¹²⁴ Patil, P.G. et al. 2016. "Toward a Blue Economy: a promise for sustainable growth in the Caribbean; an overview." The World Bank, Washington D.C. <http://documents.worldbank.org/curated/en/965641473449861013/main-report>. Accessed 2017 February

- Policies and incentives to promote and facilitate investments in blue economy
- Growth sectors and emerging market and investment opportunities

Further research by PEMSEA Country Partners, Non-Country Partners and collaborating organizations is critical for integrating the science, policy and economic

perspectives toward an understanding of blue carbon ecosystems' strategic value to the region. This report, along with the forthcoming State of Oceans and Coasts reports, can provide the foundation and a catalyst for additional, focused blue economy research to identify the interdependencies, risks and opportunities that blue carbon ecosystems/natural capital present for blue economy industries in the region.



Photo by SCA/Crooks

7 Appendices

Appendix A: Definitions

Blue carbon – The carbon stored in marine ecosystems. Blue carbon is subdivided into oceanic and coastal blue carbon component. Within this report the term blue carbon is applied only to coastal blue carbon.

Blue carbon ecosystem – A terms commonly applied to coastal vegetated ecosystems (typically mangroves, tidal marshes and seagrass meadows, but may include algal flats in arid settings) that extract carbon dioxide from the atmospheric or coastal waters and store carbon stock in biomass and soil carbon pools.

Blue economy – A practical ocean-based economic model using green infrastructure and technologies, innovative financing mechanisms and proactive institutional arrangements for meeting the twin goals of protecting our oceans and coasts and enhancing its potential contribution to sustainable development, including improving human well-being, and reducing environmental risks and ecological scarcities.

Carbon sequestration – the process by which carbon is removed directly from the atmosphere, or indirectly from coastal waters, and stored within biomass and soils.

Carbon pool – An accumulation of carbon within an ecosystems component, there are five pools measured in

carbon stock assessments: biomass (above ground, below ground and dead) litter material and soils.

Carbon stock – The amount of carbon in an ecosystem (total carbon stock) or within an individual pool (above-ground biomass, belowground biomass, soil, dead wood and litter).

Coastal blue carbon – the carbon stored in mangroves, tidal marshes and seagrass meadows.

Coastal wetland – A wetland found in coastal settings, which may be vegetated or not vegetated, connected or not connected to tides.

Mangrove – A tree, shrub, palm or ground fern that normally grows above mean sea level within the intertidal zone of marine coastal environments.

Marsh – A wetland that is dominated by herbaceous rather than woody plant species.

Salt marsh – A marsh dominated by salt-tolerant plants, such as herbs, grasses and low shrubs. Salt marshes may be terrestrial (e.g., bordering desert salt lakes) or coastal. Coastal salt marshes are found at elevations above mean tide elevation up to the highest elevation of tides.

Seagrass meadows – Seagrasses are flowering plants belonging to four plant families, all of the order Alismatales, which grow in marine, saline environments.


Tidal marsh – A marsh occurring within elevations flooded and drained by tides. May include tidal freshwater marshes at the landward extent of tides, as well as tidal brackish and salt marshes.

Tidal wetland – A wetland that lies at or beneath tidal waters, including but not limited to tidal marshes, mangroves, seagrasses as well as tidal flats.


Wetland – A land area that is saturated with water, either permanently or seasonally, such as it takes on the characteristics of a distinct ecosystem. The primary factor that distinguishes wetlands from other land forms or water bodies is the characteristic vegetation of aquatic plants, adapted to the unique hydric soil.

Appendix B: Maps of Mangrove Extent and Tidal Wetland Losses


Appendix Figure 1. Distribution of mangroves based on the 'World Atlas of Mangroves' and lands converted from tidal wetlands in Brunei Darussalam.




Appendix Figure 2. Distribution of mangroves based on the 'World Atlas of Mangroves' and lands converted from tidal wetlands in Cambodia.




Appendix Figure 3. Distribution of mangroves based on the 'World Atlas of Mangroves' and lands converted from tidal wetlands in China.




Appendix Figure 4. Distribution of mangroves based on the 'World Atlas of Mangroves' and lands converted from tidal wetlands in Indonesia.




Appendix Figure 5. Distribution of mangroves based on the 'World Atlas of Mangroves' and lands converted from tidal wetlands in Japan.



Appendix Figure 6. Distribution of mangroves based on the 'World Atlas of Mangroves' and lands converted from tidal wetlands in Malaysia.




Appendix Figure 7. Distribution of mangroves based on the 'World Atlas of Mangroves' and lands converted from tidal wetlands in Philippines.




Appendix Figure 8. Distribution of mangroves based on the 'World Atlas of Mangroves' and lands converted from tidal wetlands in Singapore.






Appendix Figure 9. Distribution of mangroves based on the 'World Atlas of Mangroves' and lands converted from tidal wetlands in Thailand.

Appendix Figure 10. Distribution of mangroves based on the 'World Atlas of Mangroves' and lands converted from tidal wetlands in Timor-Leste.




Appendix Figure 11. Distribution of mangroves based on the 'World Atlas of Mangroves' and lands converted from tidal wetlands in Viet Nam.




Appendix Figure 12. Distribution of lands converted from tidal wetlands in China.


Appendix Figure 13. Distribution of lands converted from tidal wetlands in DPR Korea.



Appendix Figure 14. Distribution of lands converted from tidal wetlands in Japan.



Appendix Figure 15. Distribution of lands converted from tidal wetlands in RO Korea.



Appendix C: Bibliography

- Alongi, D. et al. 2015. "Indonesia's blue carbon: a globally significant and vulnerable sink for seagrass and mangrove carbon." *Wetlands Ecology and Management*, 24:3-13.
- Arnell, N.W. and Gosling, S. N. 2016. "The impact of climate change on river flood risk at the global scale." *Climate Change*, 134(1):387-401.
- Association GERES. 2017. "NAMA on sustainable charcoal value-chains in Cambodia." <http://www.geres.eu/en/our-actions/item/481-nama-on-sustainable-charcoal-value-chains-in-cambodia>. <http://www.geres.eu/en/our-actions/item/481-nama-on-sustainable-charcoal-value-chains-in-cambodia>. Accessed 2017 January 25.
- Bosquet, Michelle et al. 2016. "Status Report on Nationally Appropriate Mitigation Actions (NAMAs): Mid-Year Update Report 2016." Edited by Charlotte Cuntz et al. <https://www.international-climate-initiative.com/fileadmin/Dokumente/2016/NAMA-Status-Report-June-2016.pdf>. Accessed 2016 December 20.
- Bujang, J. et al. 2006. "Distribution and significance of seagrass ecosystems in Malaysia." *Aquatic Ecosystem Health & Management*, 9:1-14.
- Costanza, R. et al. 1997. "The value of the world's ecosystem services and natural capital." *Nature*, 387:253–260.
- Costanza, R. et al. 2014. "Changes in the global value of ecosystem services." *Global Environmental Change*, 26:152-158.
- Credit Suisse. 2014. "Conservation Finance: Moving Beyond Donor Funding Toward an Investor-driven Approach." <https://www.cbd.int/financial/privatesector/g-private-wwf.pdf>. Accessed 2017 February 22.
- Creel, Liz. 2003. "Ripple Effects: Population and Coastal Regions." <http://www.prb.org/Publications/Reports/2003/RippleEffectsPopulationandCoastalRegions.aspx>. Accessed 2016 November 3.
- Crooks, S. et al. 2011. "Mitigating climate change through restoration and management of coastal wetlands and near-shore marine ecosystems: challenges and opportunities." *Environment Department Paper 121*. World Bank, Washington, DC.
- Duarte, C.M. et al. 2005. "Major role of marine vegetation on the oceanic carbon cycle." *Biogeosciences*, 2:1–8.
- FAO. 2015. "Learning Tool on Nationally Appropriate Mitigation Actions (NAMAs) in the agriculture, forestry and other land use (AFOLU) sector." <http://www.fao.org/3/a-i4642e.pdf>. Accessed 2016 December 20.
- Fortes, M. 1990. *Seagrasses: a resource unknown in the ASEAN region*. Vol. 5. WorldFish.
- Fourqurean, J.W. et al. 2012. "Seagrass ecosystems as a globally significant carbon stock." *Nature Geoscience*, 5:505-509.
- GEF Blue Forests. 2016. <http://www.gefblueforests.org/>.
- Giri, C. et al. 2011. "Status and distribution of mangrove forests of the world using earth observation satellite data." *Global Ecology and Biogeochemistry*, 20: 154-159.
- Green, E. and F. Short. 2003. *World atlas of seagrasses*. Prepared by the UIMEP World Conservation Monitoring Centre. University of California Press, Berkeley, USA.
- Hamilton, S. and D. Casey. 2016. "Creation of a high spatio-temporal resolution global database of continuous mangrove forest cover for the 21st century (CGMFC-21)." *Global Ecology and Biogeochemistry*, 25: 729-738.

- Hamilton, S. E. and D. Friess. 2016. "High-resolution annual maps of whole system mangrove carbon stocks from 2000 to 2012." *arXiv preprint arXiv:1611.00307*.
- Hansen, M. et al. 2013. "High-resolution global maps of 21st century forest cover change." *Science*, 342:850-853.
- Herr, D. and E. Landis. 2016. "Coastal blue carbon ecosystems. Opportunities for Nationally Determined Contributions." https://thought-leadership-production.s3.amazonaws.com/2016/11/12/10/26/14/0d574e9d-3eac-4591-9b8b-57001631b935/BC%20NDCs_FINAL.pdf. Accessed 2017 February 22.
- Herr, D. T. et al. 2015. Coastal "blue" carbon. A revised guide to supporting coastal wetland programs and projects using climate finance and other financial mechanisms. IUCN, Gland. https://www.iucn.org/sites/dev/files/import/downloads/wetlands_carbon_finance_paper_final_lr.pdf. Accessed 2017 January 25.
- Hiraishi, Takahiko, et al. 2014. *2013 supplement to the 2006 IPCC guidelines for national greenhouse gas inventories: Wetlands*. IPCC, Switzerland.
- Hoekstra, J. M. et al. 2010. *The atlas of global conservation: changes, challenges, and opportunities to make a difference*. Ed. J. L. Molnar. Berkeley: University of California Press.
- Howard, J. et al. 2017. "Clarifying the role of coastal and marine systems in climate mitigation." *Frontiers in Ecology and the Environment*, 15:42-50.
- Huang, X. et al. 2006. "Main seagrass beds and threats to their habitats in the coastal sea of South China." *Chinese Science Bulletin*, 51 Supp. II 136-142.
- Hutchinson, J. et al. 2013. "Predicting global patterns in mangrove forest biomass." *Conservation Letters*, 7:233-240.
- Hutomo, M. and M. Moosa. 2005. "Indonesian marine and coastal biodiversity: present status." *Indian Journal of Marine Sciences*, 34:88-97.
- Huwylar, F. et al. 2014. *Conservation finance: moving beyond donor funding toward an investor-driven approach*. Credit Suisse, WWF, McKinsey & Company: Zurich, Switzerland.
- ICAO. 2016. "Draft text for the report on agenda item 22 (Section on Global Market-based Measure Scheme)." http://www.icao.int/Meetings/a39/Documents/WP/wp_462_en.pdf. Accessed 2016 December 12.
- International Partnership for Blue Carbon. 2017. <http://bluecarbonpartnership.org/>.
- J.P. Morgan. "State of Private Investment in Conservation 2016 A Landscape Assessment of an Emerging Market." <https://www.cbd.int/financial/privatesector/g-private-wwf.pdf>. Accessed 2017 February 22.
- Jardine, S. and J. Siikamäki. 2014. "A global predictive model of carbon in mangrove soils." *Environmental Research Letters*, 9:104013.
- Joosten, H. et al. 2016. "Peatlands, forests and the climate architecture: setting incentives through markets and enhanced accounting." <https://www.umweltbundesamt.de/publikationen/peatlands-forests-the-climate-architecture-setting>. Accessed 2017 February 22.
- Kauffman, J. B. and D. Donato. 2012. *Protocols for the measurement, monitoring and reporting of structure, biomass and carbon stocks in mangrove forests*. (CIFOR Working Paper no. 86, p. 40p). Center for International Forestry Research (CIFOR), Bogor, Indonesia.
- La Viña, A. et al. 2016. "History and future of REDD+ in the UNFCCC: issues and challenges." In *Research Handbook on REDD+ and International Law*. Edited by C. Voigt. <http://www.e-elgar.com/shop/eep/preview/book/isbn/9781783478316/>. Accessed 2017 January 25.

- Lee, D. and T. Pistorius. 2015. "The Impacts of International REDD Finance." http://www.climateandlandusealliance.org/wp-content/uploads/2015/09/Impacts_of_International_REDD_Finance_Report_FINAL. Accessed 2016 December 15.
- Locatelli, T. et al. 2014. "Turning the tide: how blue carbon and payments for ecosystem services (PES) might help save mangrove forests." DOI 10.1007/s13280-014-0530-y. Accessed 2016 December 9.
- McLeod, Elizabeth and Rodney V. Salm. 2006. "Managing mangroves for resilience to climate change." World Conservation Union (IUCN), Gland. http://cmsdata.iucn.org/downloads/managing_mangroves_for_resilience_to_climate_change.pdf. Accessed 2016 December 8.
- Murray, B. et al. 2011. *Green payments for blue carbon: Economic incentives for protecting threatened coastal habitats*. Nicholas Institute for Environmental Policy Solutions, Report NI 11.04.
- NAMA. 2016. "Adaptation and mitigation initiatives in Philippine rice cultivation." <http://namanews.org/news/2015/03/05/climate-change-adaptation-oriented-nama-option-for-the-rice-sector-in-the-philippines/>. Accessed 2017 January 25.
- Nguyen Van Huy. [n.d.]. "Towards NAMA/MRV Readiness in Vietnam." [presentation]. https://unfccc.int/files/focus/mitigation/application/pdf/towards_nama_readiness_invietnam.pdf. Accessed 2017 January 25.
- Patil, P.G. et al. 2016. "Toward a Blue Economy: a promise for sustainable growth in the Caribbean; an overview." The World Bank, Washington D.C. <http://documents.worldbank.org/curated/en/965641473449861013/main-report>. Accessed 2017 January 22.
- Pearson, T. et al. 2016. "Guidance Document: Options for Nesting REDD+ Projects." (Fundación Natura Colombia 2016). http://www.v-c-s.org/wp-content/uploads/2016/07/Nesting-Options-1-Jul_Eng_final.pdf. Accessed 2016 December 18.
- PEMSEA. 2015. "Blue Economy for business in East Asia: towards an integrated understanding of Blue Economy." <http://www.pemsea.org/publications/reports/blue-economy-business-east-asia-towards-integrated-understanding-blue-economy>. Accessed 2017 January 25.
- Pendleton, L. et al. 2012. "Estimating global "blue carbon" emissions from conversion and degradation of vegetated coastal ecosystems." *PloS One*, 7.9:e43542.
- Pidgeon, E. 2009. "Carbon sequestration by coastal marine habitats: important missing sinks." In *The management of natural coastal carbon sinks*. Edited by Laffoley, D. and Grimsditch, G. IUCN, Gland.
- Resor, J. 2010. "Debt-for-nature swaps: a decade of experience and new directions for the future." <http://www.fao.org/docrep/w3247e/w3247e06.htm>. Accessed 2016 December 9.
- Restore America's Estuaries. 2016. "Blue Carbon Resources." <https://www.estuaries.org/bluecarbon-resources>.
- Richards, D. & D. Friess. 2016. "Rates and drivers of mangrove deforestation in Southeast Asia, 2000-2012." *PNAS*, 113:344-349.
- Rotich, B. et al. 2016. *Where land meets the sea. A global review of the governance and tenure dimensions of coastal mangrove forests*.
- Shi-Jun, Y. and C. Ji-yu. 1995. "Coastal salt marshes and mangrove swamps in China." *Chinese Journal of Oceanology and Limnology*, 13: 318-324.
- Short, F. et al. 2007. "Global seagrass distribution and diversity: a bioregional model." *Journal of Experimental Marine Biology & Ecology*, 350.1:3-20.

- Spalding, M. 2010. *World atlas of mangroves*. Routledge.
- Syvitskiet, J. et al. 2009. "Sinking Deltas Due to Human Activities. *Nature Geoscience*, 2:681-686.
- Twilley, R. et al. 1992. Carbon sinks in mangroves and their implications to carbon budget of tropical coastal ecosystems. In *Natural sinks of CO₂*. Springer, Netherlands, pp. 265-288.
- UNFCCC Secretariat. "Non-market based approaches: Technical Paper, FCCC/TP/2014/10, 24 November 2014." <http://unfccc.int/resource/docs/2014/tp/10.pdf>. Accessed 2017 February 22.
- UNFCCC. [n.d.]. "AR-AM0014: Afforestation and reforestation of degraded mangrove habitats --- Version 3.0." <https://cdm.unfccc.int/methodologies/DB/KMH608T6RL3P5XKNBQE2N359QG7KOE>. Accessed 2016 December 12.
- UNFCCC. [n.d.]. "Small-scale and low-income community-based mangrove afforestation project on tidal flats of three small islands around Batam City, Riau Islands Province, Republic of Indonesia." <https://cdm.unfccc.int/Projects/Validation/DB/Q94TKNG9V6GRFOTRKPLWDV2GHBCFWW/view.html>. Accessed 2016 December 12.
- UNFCCC. [n.d.]. "Subsidiary Body for Scientific and Technological Advice (SBSTA)." <http://unfccc.int/bodies/body/6399.php>. Accessed 2017 February 23.
- Vo, S. et al. 2013. "Status and trends in coastal habitats of the South China Sea." *Ocean & Coastal Management*, 85:153-163.
- World Economic Forum and Organization for Economic Cooperation and Development. 2015. "A how-to guide for Blended Finance: a practical guide for Development Finance and Philanthropic Funders to integrate Blended Finance best practices into their organizations." http://www3.weforum.org/docs/WEF_Blended_Finance_How_To_Guide.pdf. Accessed 2017 January 25.
- Zuo, P. et al. 2012. "Distribution of *Spartina* spp. along China's coast." *Ecological Engineering*, 40:160-166.





Partnerships in Environmental Management for the Seas of East Asia (PEMSEA) is an intergovernmental organization operating in East Asia to foster and sustain healthy and resilient oceans, coasts, communities and economies across the region. Through integrated coastal management solutions and partnerships, PEMSEA works with local and national governments, international development organizations, companies, investors and research institutions towards sustainable development of coasts and oceans in East Asia.